

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11)

**EP 1 223 219 A2**

(12)

**EUROPEAN PATENT APPLICATION**

(43) Date of publication:

17.07.2002 Bulletin 2002/29

(21) Application number: 02004252.9

(22) Date of filing: 30.10.1998

(51) Int Cl.7: **C12N 15/31**, C07K 14/39,  
C12N 15/81, C12N 1/19,  
C12P 21/02  
// (C12N15/31, C12R1:645),  
(C12N1/19, C12R1:645)

(84) Designated Contracting States:  
**DE FR GB**

(30) Priority: 31.10.1997 JP 31460897

(62) Document number(s) of the earlier application(s) in  
accordance with Art. 76 EPC:  
98950470.9 / 0 972 834

(71) Applicant: **ASAHI GLASS COMPANY LTD.**  
Tokyo 100-8405 (JP)

(72) Inventors:

- Tohda, Hideki, Asahi Glass Company Ltd.  
Yokohama-shi, Kanagawa 221-0863 (JP)

- Hama, Yuko, Asahi Glass Company Ltd.  
Yokohama-shi, Kanagawa 221-0863 (JP)
- Kumagai, Hiromichi, Asahi Glass Company Ltd.  
Yokohama-shi, Kanagawa 221-0863 (JP)

(74) Representative: **Müller-Boré & Partner**  
**Patentanwälte**  
**Grafinger Strasse 2**  
**81671 München (DE)**

Remarks:

This application was filed on 26 - 02 - 2002 as a  
divisional application to the application mentioned  
under INID code 62.

(54) **Inducible promoter and secretion signal for use in *Schizosaccharomyces pombe*, expression vector containing them and their use**

(57) The present invention relates to an inducible promoter gene for use in the fission yeast *Schizosaccharomyces pombe*, an inducible expression vector containing the inducible promoter gene, *Schizosaccharomyces pombe* transformed with the inducible expression vector and a process for producing a heterologous protein by using the *Schizosaccharomyces pombe* transformant.

The present invention also relates to a secretion signal gene for use in *Schizosaccharomyces pombe*, an expression vector containing the secretion signal, *Schizosaccharomyces pombe* transformed with the expression vector and a process for producing a heterologous protein by using the *Schizosaccharomyces pombe* transformant. These inducible promoter gene and secretion signal gene are originated from the *S. pombe* gene involved in expression of the invertase precursor.

The inducible promoter gene is originated from the region responsible for the catabolite repression of the

invertase precursor. When a *Schizosaccharomyces pombe* transformant carrying the inducible expression vector is incubated, the timing of the production of the desired heterologous protein can be controlled by regulating the expression of the heterologous protein structural gene depending on the presence or absence of a specific nutrient (especially glucose).

The secretion signal gene encodes the secretion signal for the invertase precursor. When a *Schizosaccharomyces pombe* transformant carrying an expression vector containing the secretion signal gene is incubated, a heterologous protein fused with the secretion signal is synthesized as a precursory protein in the cell, and after intracellular separation of the secretion signal, the heterologous protein is secreted from the cell. Therefore, it is possible to obtain the desired protein from the cell culture, and the desired heterologous protein can be purified easily.

EP 1 223 219 A2

## Description

[0001] The present invention relates to an inducible promoter gene and secretion signal gene for use in the fission yeast Schizosaccharomyces pombe (hereinafter referred to as S. pombe), an expression vector containing them and a process for producing a protein using them. In particular, it relates to a process for producing a desired protein wherein the S. pombe invertase promoter is used to make it possible to control the timing of the protein production by the presence or absence of a specific nutrient through regulated gene expression, and a process for secretory production of a desired protein by using the secretion signal gene for the S. pombe invertase precursor.

[0002] S. pombe, despite being a eukaryote, has been studied extensively for its high versatility in genetics, molecular biology and cellular biology as a unicellular organism (Nasim A. et al. eds., Molecular biology of the fission yeast, Academic Press, 1989). In its cultures, monosaccharides such as glucose and fructose are used as the main carbon sources. It is known that in a culture medium lacking these monosaccharides, expression of invertase, the enzyme that degrades sucrose into glucose and fructose, is induced to secure the carbon source necessary for its growth (Moreno S. et al., Arch Microbial. 142, 370, 1985).

[0003] S. pombe invertase is located on the cell surface and is a high-molecular weight glycoprotein with a molecular weight of about 205000, 67% of which is attributed to sugar chains composed of equimolar amounts of mannose and galactose residues. Molecular weight and amino acid studies of the protein moiety of the pure enzyme and experiments using antibodies have shown high similarity between S. pombe invertase and the invertase from the budding yeast Saccharomyces cerevisiae from the viewpoint of protein chemistry (Moreno S. et al., Biochem. J. 267, 697, 1990). It is also known that a drop in glucose concentration de-represses synthesis of invertase (Mitchinson J. et al., Cell Sci 5. 373. 1969).

[0004] Induced invertase synthesis (de-repression) is also observed in Saccharomyces cerevisiae. Previous detailed studies on genetic regulation of invertase expression, the biosynthetic pathway and the structure of the sugar chain moiety have shown that Saccharomyces cerevisiae invertase is encoded by six overlapping genes, SUC1 to SUC5 and SUC7, on one chromosome and that activation of at least one of these SUC genes leads to utilization of sucrose and raffinose (Hohmann S. et al., Curr Genet 11, 217, 1986).

[0005] In contrast, with respect to S. pombe, although purification of the invertase protein has been reported (Moreno S. et al. 1985), no invertase genes had been identified until the present inventors and coworkers recently reported two overlapping invertase genes *inv0+* and *inv1+* in S. pombe. Because *inv0+* is likely a pseudogene having an incomplete open reading frame, *inv1+* is the only one gene encoding S. pombe invertase, which is supposed to confer the ability to grow on sucrose even in the absence of other carbon sources ("Kobogaku" edited by Yositaka Hashitani, Iwanam Shoten. 1967).

[0006] Analysis of the promoter region of the isolated gene suggested that a specific sequence between the 1st and 62nd base pairs is involved in catabolite repression.

[0007] In Saccharomyces cerevisiae, the SUC2 gene is transcribed into two messenger RNAs (mRNAs) from different transcription initiation sites. The shorter one is a constitutive mRNA encoding the intracellular invertase, while the longer one is a mRNA encoding the catabolite-repressible secretory invertase with a de-repression ratio of not less than 200 (Carlson M. et al., Mol. Cell. Biol. 3, 439, 1983). Analysis of the promoter region for the longer mRNA suggested that the transcription initiation factor binds to a specific repeated sequence between positions -650 and -418 (Salokin L et al., Mol. Cell. Biol. 6, 2314, 1986). The region between positions -418 and -140 has been shown to be necessary for glucose repression.

[0008] These regions in the SUC2 gene showed no significant homology with the *inv1+* upstream region between positions 1 and 2809. However, multiple copies of a so-called 7-bp motif with the sequence (A/C) (A/G)GAAAT, which is repeated at five sites in the region indispensable for glucose derepression, have been found in the *inv1+* upstream region. Further, while palindrome stem-loops have been identified at almost the same positions in the upstream regions of glucose-repressible genes (SUC, MAL and GAL), palindrome sequences have also been found in the upstream region of the *inv1+* gene from S. pombe. These sequences are anticipated to play an important role in glucose repression in S. pombe.

[0009] The yeast S. pombe is phylogenetically different from Saccharomyces cerevisiae. It is quite different from other yeasts in the chromosome structure and various mechanisms for genome replication, RNA splicing, transcription and posttranslational modification, and rather resembles animal cells in some of these aspects. For this reason, S. pombe is widely used as a eukaryotic model (Giga-Hama and Kumagai, eds., Foreign gene expression in fission yeast Schizosaccharomyces pombe, Springer-Verlag, 1997).

[0010] S. pombe is also widely used as a host for expression of heterologous protein genes and known to be suited especially for expression of genes from animals including human (JP-A-5-15380 and JP-A-7-163373). For its advanced membrane structures including the Golgi body and the endoplasmic reticulum, S. pombe is also used for expression of membrane proteins and shows high level expression. For S. pombe, constitutive expression vectors (pEVP11, pART1 and pTL2M) and an inducible expression vector using the promoter region of the *nmt1+* gene (pREP1) are usually

used as expression vectors. No S. pombe expression vectors of the GAL type or the SUC type have been known though these types of vectors are commonly used for Saccharomyces cerevisiae.

[0011] The expression of the SUC2 gene from Saccharomyces cerevisiae in S. pombe has been shown to be constitutive, not catabolite repressible, though the expression product contains galactose residues conferred by the host (Zarate, V. et al., J Applied Bacteriology, 80, 45, 1996), suggesting differences between S. pombe and Saccharomyces cerevisiae in the mechanism for catabolite repression of invertase. The differences are of great significance because the promoter from Saccharomyces cerevisiae usually used by those skilled in the art for construction of inducible expression vectors of the invertase type (the SUC2 type) is not applicable to S. pombe vectors. Therefore, development of S. pombe vectors of this type has been long delayed.

[0012] On the other hand, the present inventors constructed an expression vector using the secretion signal gene encoding the secretion signal in the precursor of a S. pombe mating pheromone (WO96/23890). However, this secretion signal gene is not an all-purpose secretion signal gene, and other secretion signal genes that function in S. pombe are desired for production of some types of protein.

[0013] As a result of their extensive research with a view to solving the above problems, the present inventors have accomplished the present invention by preparing a new clone of the S. pombe invertase gene and constructing an inducible expression vector. They have also found that the N-terminal 22 amino acid sequence in the amino acid of the invertase precursor functions as a secretion signal. On the basis of these findings, they have constructed an expression vector using the secretion signal gene and established secretory production of desired proteins.

[0014] The present invention relates to a region in the invertase gene from S. pombe, which is involved in catabolite repression, an inducible expression vector using the region and a system using it for heterologous gene expression and provides:

a DNA in an invertase gene from Schizosaccharomyces pombe, which is located in a region involved in catabolite repression,

a DNA having the base sequence of bases 1 to 2809 in SEQ ID NO: 1 in the Sequence Listing,

a recombinant vector containing the sequence of the DNA, a multicloning vector containing the sequence of the DNA and a multicloning site,

a multicloning vector having the structure shown in Figure 9,

an expression vector for transformation of Schizosaccharomyces pombe containing the sequence of the DNA and

a heterologous protein structural gene,

a transformant from Schizosaccharomyces pombe containing the expression vector, and

a process for producing a protein which comprises incubating the transformant and recovering an expressed heterologous protein.

[0015] Firstly, the present inventors cloned and sequenced a S. pombe invertase gene, which had not been genetically identified. Then, they demonstrated by gene disruption analysis that the invertase gene is responsible for the overall invertase activity. Further, they identified the region involved in catabolite repression and constructed an inducible expression vector using the region. They actually constructed a recombinant vector carrying the gene of a green fluorescent protein, transformed S. pombe with the vector and confirmed the expression of the protein by assay of invertase activity and immunological analysis. They also demonstrated repression of the heterologous gene expression in the presence of glucose in the culture medium and derepression by exhaustion of glucose.

[0016] The present inventors used the following procedure to identify and characterize the gene of the S. pombe invertase precursor:

(1) PCR using a cDNA library from S. pombe as a template and primers based on conserved amino acid sequences in invertase genes from many other organisms;

(2) screening of a genomic library from S. pombe by plaque hybridization using the PCR product as a probe for positive clones;

(3) confirmation of the positive clones by restriction digestion followed by electrophoresis;

(4) Southern hybridization analysis and total sequencing of a fragment with a specific length in the positive clones;

(5) gene disruption analysis of invertase activity;

(6) investigation of the optimum pH for expression of the invertase gene from S. pombe and the effects of the glucose concentration on glucose repression and derepression; and

(7) identification of a region indispensable for glucose repression through subcloning of the related upstream region.

[0017] Also, the present inventors constructed a S. pombe invertase inducible expression vector by the following procedure and actually demonstrated inducible expression of a green fluorescent protein:

- (1) construction of an inducible multicloning expression vector pRI0M containing an invertase promoter by modifying a S. pombe multicloning vector, pTL2M (JP-A-7-163373);
- (2) construction of an inducible expression vector pRI0EGFP for expression of a green fluorescent protein variant from the inducible multicloning vector, pRI0M;
- (3) transformation of a wild-type S. pombe strain with the inducible expression vector, pRI0EGFP, for expression of the green fluorescent protein variant;
- (4) demonstration of the expression of the green fluorescent protein variant by activity (fluorescence) analysis and SDS-PAGE-western blotting; and
- (5) establishment of the conditions for the inducible expression on the basis of the dependence of the expression level on the glucose concentration in the culture medium.

[0018] SEQ ID NO: 1 in the Sequence Listing is the base sequence of the gene of the invertase precursor, which contains a region involved in catabolite repression. The region involved in catabolite repression is the DNA sequence between positions 1 to 2809 or within the DNA sequence. In the DNA sequence between positions 1 and 2809, the region extending from position 1 to position 620 and the region extending from position 1610 to position 2610 are especially important, as is evident from the results of the analysis in Example 6 shown in Figure 8 (position 2810 in SEQ ID NO: 1 corresponds to position 1 in Figure 8). This means that the inducible promoter in the present invention is not restricted to a DNA having the base sequence from position 1 to position 2809 so long as it contains these genes involved in catabolite repression and functions as an inducible promoter. Still, a DNA having a base sequence from position 1 to position 2809 is preferred as an inducible promoter because it actually functions in S. pombe.

[0019] The above-mentioned DNA which contains genes involved in catabolite repression and function as an inducible promoter, preferably having the base sequence from position 1 to position 2809, is hereinafter referred to as the inducible promoter gene. The inducible promoter gene can be integrated with a vector for construction of recombinant vectors such as multicloning vectors and expression vectors. A multicloning vector is a vector having a multicloning site and provides an expression vector through introduction of a desired structural gene into the multicloning site. An expression vector is a vector containing a structural gene and used for expression of a structural gene encoding a heterologous protein. A "heterologous" protein is a protein which is not inherent in the host. For example, when the host is S. pombe, a heterologous protein is a protein which is not inherent in S. pombe (such as a human protein).

[0020] In the expression vector, the inducible promoter gene is located upstream from the heterologous protein structural gene and regulates expression of the structural gene. The inducible promoter gene in the expression vector regulates the expression of the heterologous protein structural gene downstream, like the inducible promoter gene, located upstream in the base sequence represented by SEQ ID NO: 1 regulates the expression of the structural gene of the invertase precursor. In the multicloning vector, the inducible promoter gene is located upstream from the multicloning site into which a heterologous protein structural gene is to be introduced.

[0021] One example of the multicloning vector of the present invention is the multicloning vector pRI0M constructed in Example 7 and has the structure shown in Figure 9. The entire base sequence of pRI0M is SEQ ID NO: 2. Inv1-P is the above-mentioned inducible promoter gene, and MCS is the multicloning site. One example of the expression vector of the present invention is the inducible expression vector pRI0EGFP for expression of a green fluorescent protein variant constructed in Example 8 by introducing the structural gene (EGFP-ORF) of a green fluorescent protein variant and has the structure shown in Figure 10. The entire base sequence of the expression vector pRI0EGFP is SEQ ID NO: 13.

[0022] The most suitable cell (host) to transform with the expression vector of the present invention is S. pombe because the inducible promoter gene in the present invention is an inducible promoter gene from S. pombe.

[0023] Under catabolite repressing conditions (for example, in a culture medium containing a high level of glucose), S. pombe transformed with the expression vector of the present invention grows with no (or low) expression of the heterologous protein. Growth at this stage without the burden of heterologous protein expression is more efficient than growth under the burden. Subsequent incubation under catabolite derepressing conditions (for example, in a culture medium containing no or a low level of glucose) invites the increased number of S. pombe cells to high level expression of the heterologous protein, though growth of S. pombe is less efficient than under catabolite repressing conditions. Thus, controlled transition between growth of S. pombe and heterologous protein expression through catabolite repression allows more efficient production of a heterologous protein.

[0024] Catabolite repression can be controlled not only in an active way as described above but also in a passive way. For example, when a S. pombe transformant is incubated in a culture medium containing a given amount of glucose, the S. pombe grows under catabolite repressing conditions containing a high level of glucose in the initial stage, but later on production of a heterologous protein predominates due to catabolite derepression as glucose is exhausted. This way, more efficient heterologous protein production of than ever is possible without active control of the glucose level.

[0025] In addition to the above-mentioned total sequencing of the invertase precursor gene from S. pombe, the

present inventors determined the complete amino acid sequence of the invertase precursor (the amino acid sequence in SEQ ID NO: 1). Then, they have found that the first 22 amino acid peptide in the amino acid sequence of the invertase precursor (Met Phe Leu Lys Tyr Ile Leu Ala Ser Gly Ile Cys Leu Val Ser Leu Leu Ser Ser Thr Asn Ala) acts as a secretion signal. Hereinafter, the peptide is referred to as the secretion signal.

**[0026]** It is expected that a desired heterologous protein produced in transformed *S. pombe* cells as a protein fusion having the secretion signal at the N-terminal is secreted from the cells after intracellular processing which splits the protein fusion into the secretion signal and the heterologous protein. The present inventors constructed an expression vector carrying a heterologous protein structural gene (specifically, human interleukin 6-a'c1 variant) fused with a DNA encoding the secretion signal (namely, a structural gene of a protein fusion as mentioned above) and demonstrated secretion of the heterologous protein from *S. pombe* cells transformed with the expression vector.

**[0027]** The present invention provides the secretion signal, a DNA encoding the secretion signal (hereinafter referred to as a secretion signal gene), a recombinant vector carrying the secretion signal gene, a multicloning vector carrying the secretion signal gene, an expression vector carrying the secretion signal gene and a heterologous protein structural gene for transformation of *S. pombe*, a *S. pombe* transformant carrying the expression vector and a process for producing a protein which comprises incubating the transformant and recovering the expressed heterologous protein.

**[0028]** The secretion signal gene is not restricted to the 66-bp sequence extending from position 2810 to position 2875 in SEQ ID NO: 1 and may be a DNA having a different base sequence encoding the amino acid sequence of the secretion signal. In the expression vector, the secretion signal and the heterologous protein structural gene is preferably linked directly. But they may be linked via another DNA sequence, for example, extending from position 2876 in SEQ ID NO: 1. In this case, the protein product has extra amino acid residues at the N-terminal of the heterologous protein but can be converted into the desired heterologous protein by trimming off the N-terminal extra amino acid residues. However, the disadvantage from the presence of these extra amino acid residues usually becomes more serious for the desired protein as the number of extra amino acid residues increases. Therefore, as the intervening DNA between the secretion signal gene and the heterologous protein structural gene, a short DNA encoding at most 10 amino acid residues is preferable. Particular preferably, the secretion signal gene and the heterologous protein structural gene are linked directly.

**[0029]** Construction of an expression vector carrying the secretion signal gene using a multicloning vector can be attained by inserting a heterologous protein structural gene fused with the secretion signal gene into the multicloning site of a multicloning vector or by inserting a heterologous protein structural gene into the multicloning site of a multicloning vector carrying the secretion signal gene. The latter method tends to restrict the structure of the multicloning site because the secretion signal gene is preferred to be located immediately in front of the multicloning site as described above. Therefore, the former method is preferred for construction of an expression vector. As a *S. pombe* multicloning vector, for example, pTL2M, which is disclosed in JP-A-163373, is preferable.

**[0030]** According to the present invention, an expression vector can be constructed by using both the inducible promoter gene and the secretion signal gene. For example, an expression vector which contains the DNA sequence of from position 1 to position 2875 in SEQ ID No: 1 and a heterologous protein structural gene introduced downstream of the DNA sequence can be constructed. Such an expression vector enables catabolite repressible secretory production of a heterologous protein by the host cell. A similar expression vector can be constructed by using a known secretion signal gene (such as the secretion signal gene disclosed in WO96/23890) instead of the above-mentioned secretion signal gene.

**[0031]** Figure 1 shows a comparison of (partial) amino acid sequences deduced from *inv1*<sup>+</sup>, the *Schwanniomyces occidentalis* invertase gene and the fission yeast *SUC2* gene. Figure 2 is the restriction map of the *inv1*<sup>+</sup> gene. Figure 3 electrophoretically shows disruption of the *inv1*<sup>+</sup> gene. Figure 4(a) is a photograph of colony gel overlay assay of invertase activity (for phenotype characterization) as a substitute for a drawing. Figure 4(b) is a schematic explanation of experimental design of the invertase activity assay shown in Figure 4(a). Figure 5(a) is a photograph of colony gel overlay assay of invertase activity (for phenotype characterization) as a substitute for a drawing. Figure 5(b) is a schematic explanation of experimental design of the invertase activity assay shown in Figure 5(a).

**[0032]** Figure 6 graphically shows the relation between the invertase activity and the glucose concentration. Figure 7 graphically shows the relation between the invertase activity and the glucose concentration. Figure 8 shows the results of analysis of invertase promoters. Figure 9 show the structure of an inducible expression vector pRIOM. Figure 10 shows the structure of an inducible expression vector pRI0EGFP for expression of a green fluorescent protein.

**[0033]** Figures 11(a) and 11(b) demonstrate the expression of a green fluorescent protein. Figures 12(a) and 12(b) graphically show the relation between the incubation time and the expression level of the green fluorescent protein. Figures 13(a) and 13(b) graphically show the relation between the incubation time and the expression level of the green fluorescent protein. Figures 14(a) and 14(b) show the relation between the incubation time and the expression level of the green fluorescent protein. Figure 15 is a SDS-PAGE pattern obtained in analysis of the expression of interleukin 6a'c1 variant. Figure 16 is the western blot pattern of the expressed interleukin variant.

**[0034]** Now, the present invention will be described in further detail with reference to specific Examples.

## EXAMPLE 1

Isolation of *S. pombe* invertase gene

[0035] PCR using a *S. pombe* cDNA library as the template and primers designed on the basis of conserved sequences in invertase genes from other organisms shown in SEQ ID NOS: 3 to 5, gave amplification products of about 300bp and about 400bp. Each PCR product was purified by using EASY TRAP (Takara Shuzo Co., Ltd.) and sequenced after ligation into a vector by using pMOS Blue T vector kit (Amersham Pharmacia Biotech K.K.). The deduced amino acid sequence indicated that part of the 400-bp PCR product has significant homology with the SUC2 gene from

*Saccharomyces cerevisiae*.

[0036] Screening of a genomic library of *S. pombe* for the entire invertase gene by plaque hybridization using the 400-bp PCR product as a probe picked up 15 positive clones from about 8,000 plaques. Secondary screening of the positive clones by plaque hybridization left four positive clones. Treatment of small amounts of phage DNA extracts from the positive clones with various restriction enzymes gave identical cleavage patterns, and thus revealed that all the clones were identical.

[0037] The entire invertase gene was isolated by the following two-step procedure. A 3.0-kb HindIII fragment was isolated from the hybrid-forming clones and ligated into a plasmid pBluescript II SK- (Toyobo Co., Ltd.). Restriction mapping identified BamHI and Sall sites. Subcloning of the fragment using these restriction enzymes and subsequent sequencing using a deletion technique revealed that the HindIII fragment contains the complete ORF of the gene but contains only about 200 bp within the upstream region, which is supposedly involved in the gene expression. Therefore, separately, a 3.5-kb BamHI fragment from the hybrid-forming clones was further digested with HindIII to give a 2.6-kb fragment. Subcloning of the 2.6-kb fragment in plasmid pBluescript II SK-, and subsequent sequencing using a deletion technique revealed that the BamHI-HindIII fragment contained a sequence within the upstream region supposedly involved in the gene expression. The resulting complete 5.6-kb gene was designated as *inv1+*. The base sequence of *inv1+* and the amino acid sequence encoded by its ORF are shown in SEQ ID NO: 1. A plasmid carrying the complete gene was designated as pINV3000.

[0038] These results suggest that the *inv1+* product has 16 asparagine-linked glycosylation sites. Figure 1 shows a comparison of the amino acid sequence deduced from the base sequence of the *inv1+* gene from *S. pombe*, the amino acid sequence of *Schwanniomyces occidentalis* invertase and the amino acid sequence deduced from the base sequence of the SUC2 gene from *Saccharomyces cerevisiae*. Amino acids that are common to the three are marked with \*. Figure 1 clearly shows the amino acid sequence deduced from the *inv1+* base sequence has significant homology with invertases from other origins such as *Schwanniomyces occidentalis* and *Saccharomyces cerevisiae*, which suggests the *inv1+* encodes invertase.

## EXAMPLE 2

Disruption of the *inv1+* gene

[0039] The HindIII site in plasmid pBluescript II SK- having a *S. pombe* *ura4+* gene insert at the ClaI site was disrupted by HindIII digestion followed by blunting and self-ligation (self-cyclization). Double restriction digestion of the plasmid with XbaI and HincII gave a *ura4+* fragment. The plasmid pBluescript was integrated with the *ura4+* fragment after restriction digestion with SpeI and subsequent blunting and XbaI digestion, to provide a plasmid having BamHI sites on both sides of *ura4+*. The BamHI site in plasmid pBluescript II SK- was disrupted similarly by restriction digestion followed by blunting and self-ligation, and the 3.0-kb fragment containing the *inv1+* ORF was inserted at the HindIII site. BamHI digestion of the plasmid eliminated a 1.4-kb fragment (containing part of the *inv1+* ORF encoding the C-terminal of invertase) from the 3.0-kb insert, and a *ura4+* cassette having BamHI sites at both ends was inserted. HindIII digestion of the resulting plasmid gave a DNA fragment having *inv1+* neighboring regions at both ends (Figure 2). The restriction map of the *inv1+* gene is shown in Figure 2, wherein the open reading frame (ORF) is indicated by the arrow (*inv1+*ORF) and the *ura4+* replacement from *Schizosaccharomyces pombe* is boxed (*ura4+*). The disruption mutant strain had an *inv1+* fragment carrying the *S. pombe* *ura4+* gene instead of the 1.4-kb *inv1+* BamHI-BamHI fragment partly containing the ORF. The *inv1+* fragment was used to transform a wild-type *S. pombe* strain, TP4-1D [*h<sup>-</sup>, leu1, ura4, ade6-M216, his2*, obtained from Dr. Takashi Toda (Imperial Cancer Research Foundation)], and viable colonies on a uracil-free culture medium were collected. Overlay assay of invertase activity revealed that 7 out of 28 strains, namely 25% of the *ura4+* colonies, lacked invertase activity.

[0040] Further, to verify the chromosomal *inv1+* gene disruption, genomic DNA from a strain lacking invertase activity was analyzed after double restriction digestion with HindIII and Sall by Southern hybridization using the *inv1+* HindIII-Sall fragment (2 kb) as the probe. The 3-kb hybridized fragment, which was not digested with Sall, shown in Figure 3 demonstrates that part of the *inv1+* gene in the chromosomal DNA had been replaced with the *ura4+* gene in the

strain which lacked invertase activity.

[0041] Thus, the *inv1<sup>+</sup>* gene proved to be the only one invertase gene expressed in *S. pombe*.

### EXAMPLE 3

#### Restoration of invertase activity by the *inv1<sup>+</sup>* gene

[0042] The 3.0-kb HindIII fragment containing the entire *inv1<sup>+</sup>* ORF from the invertase gene was inserted into *S. pombe* vector pAU-SK (obtained from Dr. Chikashi Shimoda, Department of Science, Osaka City University), and the resulting recombinant vector was used to transform the invertase-defective strains (Example 2). The resulting transformants were streaked on YP sucrose plates (supplemented with 10 µg/ml antimycin A and 20 µg/ml bromocresol purple) for overlay assay of invertase activity. Further, the 2.6-kb *inv1<sup>+</sup>* BamHI-HindIII fragment containing the upstream promoter region and the 2.0-kb HindIII-Sall fragment from the invertase gene containing the ORF were ligated, and the resulting 4.6-kb BamHI-Sall fragment was inserted into pAU-SK. Transformation of the resulting recombinant vector into the invertase-defective strain was followed by similar overlay assay of invertase activity.

[0043] Both transformants (*inv1Δ*[pAU-SK::*inv1<sup>+</sup>*]) and the wild-type strain TP4-1D (WT) developed blue stains, which indicate invertase production, unlike the *inv1<sup>+</sup>* disruption mutant strain (*inv1Δ*). The addition of the upstream promoter region resulted in stronger stains, which suggest high-level invertase expression (Figure 4(a) and 4(b)). Figure 4(a) is a photograph showing the results of the gel overlay assays, and Figure 4(b) is a schematic explanation of the stained sections shown in Figure 4(a).

[0044] The invertase-defective strain were hardly viable on YP sucrose plates (supplemented with 10 µg/ml antimycin) whereas the wild-type strain and the transformants were recognizably viable after 5 days incubation at 30°C (Figure 5(a) and (b)). Figure 5(a) is a photograph showing the results of characterization by colony formation, and Figure 5(b) schematically explains the characterization shown in Figure 5(a).

[0045] These results demonstrate that the *inv1<sup>+</sup>* gene expression product is the invertase located on the cell surface.

### EXAMPLE 4

#### Determination of glucose concentration for gene repression

[0046] For determination of the critical glucose concentration for catabolite repression of the invertase gene, the wild-type strain TP4-1D was incubated at 30°C in 5 ml of MM medium containing 2%, 4%, 8% and 16% glucose with shaking to the mid-logarithmic growth phase. Invertase assays were done by the method of Goldstein et al., and the post-incubational glucose concentrations in the medium were determined by the phenol-sulfate method (Figure 6). The hatched bars indicate invertase activity per cell (U/OD), and the empty bars indicate the residual glucose concentration. Judging from the graph, a glucose concentration of 8% is the optimum for glucose repressing incubation, because when the glucose concentration was 8%, the invertase activity was sufficiently repressed with little decrease of glucose.

### EXAMPLE 5

#### Determination of glucose concentration for induced invertase production

[0047] For determination of the most effective glucose concentration for induced invertase production, the wild-type strain TP4-1D and a transformant [obtained by transforming the invertase-defective strain (Example 3) with a pAU-SK vector carrying the *inv1<sup>+</sup>* BamHI-Sall fragment] were preincubated in a medium containing 2% glucose to the mid-logarithmic growth phase and incubated in an MM medium containing 0%, 0.01%, 0.05%, 0.1%, 0.25%, 0.5%, 1.0% and 2% glucose with shaking at 27°C for 3 hours, and the invertase activity was assayed (Figure 7). Each run of assay was carried out at 30°C over 180 minutes. 1U of invertase converts 1 nmol of sucrose into glucose per 1 minute at 30°C, pH 4.0.

[0048] The optimum glucose concentration for induction was found to be 0.1% for the wild-type strain and 0.05% for the transformant. The invertase activity in the wild-type strain was 40 times higher under derepressing conditions than under repressing conditions. These results demonstrate catabolite repression in *S. pombe*.



## EXAMPLE 6

Analysis of the inv1<sup>+</sup> promoter region

5 [0049] Fragments obtained ligating *S. pombe* inv1<sup>+</sup> upstream sequences extending from positions 1, 620, 1100, 1610 and 2610, respectively, and the inv1<sup>+</sup> ORF were inserted into expression vector pAU-SK to obtain 5 plasmids for deletion studies. The plasmids containing the upstream sequences extending from positions 1, 1610 and 2610 carry the inv1<sup>+</sup> BamHI-Sall, SacI-Sall and HindIII-Sall fragments, respectively. The plasmids containing the inv1<sup>+</sup> upstream sequences extending from positions 620 and 1100 were constructed by site specific introduction of a SpeI site into

10 pAU-SK::inv1<sup>+</sup> (BamHI-Sall) using primers shown in SEQ ID NOS: 6 and 7, respectively, followed by partial removal of the upstream region by SpeI treatment.

[0050] The plasmids thus obtained were used to transform the invertase-defective strain (Example 3). Invertase assays were done to determine the enzyme activity in each transformant (Fig. 8). The results suggest that the sequence between position 1 and position 620 is essential for glucose repression. The region between position 1620 and position

15 2610 was identified as essential for high-level glucose derepression of invertase.

## EXAMPLE 7

Construction of inducible multicloning expression vector pRI0M carrying the invertase promoter

20 [0051] PCR amplification using the plasmid pINV3000 (Example 1) carrying the invertase gene from *S. pombe* as the template and oligo DNAs shown in SEQ ID NOS: 8 and 9 as the primers was performed to give a sequence which contains the promoter region for the invertase gene and has restriction enzyme recognition sequences at both ends. After terminal double restriction digestion with SpeI (Takara Shuzo Co., Ltd.) and EcoRI (Takara Shuzo Co., Ltd.), the

25 sequence was subjected to agarose gel electrophoresis. Purification of the band of about 3000 bp by the glass beads method using EASY-TRAP gave an insert fragment.

[0052] The *S. pombe* multicloning vector pTL2M (JP-A-7-163373) was subjected to agarose gel electrophoresis after terminal double restriction digestion with SpeI and EcoI. Purification of the band of about 4500 bp by the glass beads method using EASY-TRAP gave a vector fragment.

30 [0053] The two fragments were ligated with a DNA ligation kit (Takara Shuzo Co., Ltd.) and transformed into *E. coli* strain DH (Toyobo Co., Ltd.). *E. coli* colonies were screened for the inducible expression vector pRI01M shown in Figure 9 and SEQ ID NO: 2 in the Sequence Listing through base sequencing and restriction mapping, and the inducible expression vector was recovered by the alkali-SDS method on a preparatory scale.

## 35 EXAMPLE 8

Construction of inducible expression vector pRI0EGFP for expression of green fluorescent protein

40 [0054] PCR amplification using the plasmid pINV3000 (Example 1) carrying the invertase gene from *S. pombe* as the template and oligo DNAs shown in SEQ ID NOS: 8 and 10 as the primers was performed to give a sequence which contains the promoter region for the invertase gene and has restriction enzyme recognition sequences at both ends. After terminal double restriction digestion with SpeI and NheI (Takara Shuzo Co., Ltd.), the sequence was subjected to agarose gel electrophoresis. Purification of the band of about 3000 bp by the glass beads method using EASY-TRAP gave a fragment for use as the promoter insert.

45 [0055] PCR using the plasmid pEGFP carrying the jellyfish (*Aequorea victoria*) green fluorescent protein variant gene (Clontech) as the template and oligo DNAs shown in SEQ ID NOS: 11 and 12 in the Sequence Listing as the primers was performed to amplify the ORF in the green fluorescent protein variant gene. The PCR product was subjected to agarose gel electrophoresis after terminal double restriction digestion with NheI and HindIII. Purification of the band of about 700 bp by the glass beads method using EASY-TRAP gave a fragment for use as the ORF insert.

50 [0056] The *S. pombe* multicloning vector pTL2M was cleaved by double restriction digestion with SpeI and HindIII and then subjected to agarose gel electrophoresis. Purification of the band of about 4500 bp by the glass beads method using EASY-TRAP gave a vector fragment.

[0057] The three fragments were ligated with a DNA ligation kit and transformed into *E. coli* strain DH5. *E. coli* colonies were screened for the inducible expression vector pRI0EGFP for expression of the green fluorescent protein variant shown in Figure 10 and SEQ ID NO: 13 in the Sequence Listing through base sequencing and restriction mapping, and the inducible expression vector was recovered by the alkali-SDS method on a preparatory scale.

55



## EXAMPLE 9

Preparation of *S. pombe* transformant ASP138

[0058] *S. pombe* wild-type strain ARC001 [leu1-32h<sup>-</sup> (isogenic to ATCC38399)] was transformed with the inducible expression vector pRI0EGFP for expression of the green fluorescent protein variant and a transducing vector pAL7 as described by Okazaki et al. (Okazaki et al., "Nucleic Acids Res.", 18, 6485-6489, 1990).

[0059] 1 ml of a preincubated ARC001 culture in YPD medium was incubated in minimum medium MB+Leu with shaking at 30°C for 14 hours to a cell density of  $3 \times 10^7$  per 1 ml, and the cells were collected, washed with water, suspended in 1 ml of 0.1M lithium acetate (pH 5.0) and incubated at 30°C for 60 minutes. A 100 µl portion of the suspension was mixed with 4 µg of the inducible expression vector pRI0EGFP and 0.5 µg of PstI-digested pAL7 in 15 µl TE and further with 290 µl of 50% PEG 4000 thoroughly and incubated at 30°C for 60 minutes, at 43°C for 15 minutes and at room temperature for 10 minutes, successively. After centrifugal removal of PEG, the cells were suspended in 1 ml of 1/2 YEL+Leu medium. After 10-fold dilution, 1 ml of the suspension was incubated at 32°C for 2 hours, and a 300 µl portion was spread on minimum medium agar MMA. After 3 days of incubation at 32 °C, about 300 independent colonies had developed on the plate.

[0060] 10 colonies of the transformant were inoculated in 2 ml of YEL medium containing 10 µg/ml antibiotic G418 (YEL10 medium) and incubated with shaking at 32°C. 2 days later, 6 clones were viable. In their subcultures, 4 clones were viable 3 days later. The putative desired transformant (ASP138 strain) was frozen in glycerol and stored for use in subsequent experiments.

## EXAMPLE 10

Analysis of expression of green fluorescent protein variant

[0061] *S. pombe* transformant ASP138 (Example 9) was inoculated in YPD medium containing 100 µg/ml G418 (YPD100) and incubated at 32°C for 2 days. Green fluorescence was observed from each cell under a fluorescence microscope (excitation wavelength 490nm/emission wavelength 530nm). Green fluorescence emission was also observed upon ultraviolet irradiation from the centrifugally collected cells. Thus, expression of the desired green fluorescent protein in the active form was confirmed.

[0062] Strain ASP138 (Example 9) was incubated in 5 ml YPD100 at 32°C for 3 days, collected, washed and suspended in 50mM tris-HCl (pH 7.5), disrupted with glass beads in a mini bead beater (Biospec). After removal of the glass beads, the cell extract was heated in the presence of SDS (1%) at 80°C for 15 minutes. Separately, a negative control was extracted from the transformant carrying pR10M by the same procedure.

[0063] 50 µg protein from the extract was analyzed by SDS-polyacrylamide gel electrophoresis (Figure 11(a) and (b)). After Coomassie Brilliant Blue (CCB) staining, the extract and a recombinant green fluorescent protein (Clontech) as the positive control showed major bands with a molecular weight of 25000, but the negative control did not. Further, 50 µg protein from the extract was analyzed by SDS-polyacrylamide gel electrophoresis followed by western blotting on a PVDF membrane using an anti-green fluorescent protein antibody (Clontech). The cell extract and the positive control showed major bands with a molecular weight of 25000, but the negative control did not. These results provide biological evidence of expression of the desired green fluorescent protein.

## EXMAPLE 11

Optimization of the incubation method (1)

[0064] The transformed *S. pombe* strain ASP (Example 9) was incubated in YPD medium containing 100 µg/ml G418 (YPD100) at 32°C. The expression level of the green fluorescent protein in the cell culture was determined fluorometrically by means of a microplate reader (Corona Electric Co., Ltd.) equipped with a fluorescent attachment (excitation wavelength 490nm/ emission wavelength 530nm) (Figure 12(a) and (b)). OD, FLU/OD and time denote the cell density, the fluorescence intensity per cell and the incubation time, respectively. The results show that strain ASP138 did not express the green fluorescent protein variant until the late-growth phase after glucose exhaustion in the mid-growth phase, unlike strain ASP122 having a non-inducible cytomegalovirus promoter [a transformant carrying a recombination product of pHGFPs65T (Clontech) for expression of a green fluorescent protein variant of S65T prepared as disclosed in JP-A-7-163373], clearly due to repression of the inducible invertase promoter in the presence of glucose and subsequent derepression by glucose exhaustion, demonstrating the applicability of this mechanism to expression of the green fluorescent protein as a heterologous protein.

## EXAMPLE 12

Incubation of strain ASP138 (2)

[0065] The transformed *S. pombe* strain ASP138 (Example 9) was incubated in YPD medium containing 100 µg/ml G418 (glucose concentration 8%) at 32°C to the mid-growth phase, and after medium change, incubated in YPDG medium containing 100 µg/ml G418 (glucose concentration 0.1%, glycerol concentration 3%) (Figure 13(a) and (b)). OD, FLU/OD and time denote the cell density, the fluorescence intensity per cell and the incubation time, respectively. The results show that while the green fluorescent protein was not expressed in the cells incubated without medium change (Untreated), expression of the green fluorescent protein in the cells incubated with the medium change was activated by the medium change (Medium-changed) probably because the depletion of glucose in the medium provoked derepression of the invertase promoter and thereby induced the protein expression. The high level expression induced by the medium change to a low-glucose expression medium suggests that use of a growth medium (with a high glucose concentration) and an expression medium (with a low glucose concentration) can differentiate between cell growth and protein expression. It was demonstrated that the repression of the inducible invertase promoter in the presence of glucose and derepression by exhaustion of glucose could be utilized in expression of the green fluorescent protein as a heterologous protein.

## EXAMPLE 13

Incubation of strain ASP138 (3)

[0066] The transformed *S. pombe* strain ASP138 (Example 9) was incubated in YPD medium containing 100 µg/ml G418 (glucose concentration 8%) at 32°C to the late-growth phase, and after medium change, incubated in YPDG medium containing 100 µg/ml G418 (glucose concentration 0.1%, glycerol concentration 3%) (Figure 14(a) and (b)). OD, FLU/OD and time denote the cell density, the fluorescence intensity per cell and the incubation time, respectively. The results show that while the green fluorescent protein was not expressed in the cells incubated without medium change (Untreated), expression of the green fluorescent protein in the cells incubated with the medium change was activated by the medium change (Medium-changed) probably because the depletion of glucose in the medium provoked derepression of the invertase promoter and thereby induced the protein expression. The high level expression induced by the medium change to a low-glucose expression medium from a high-glucose medium before glucose exhaustion suggests that use of a growth medium (with a high glucose concentration) and an expression medium (with a low glucose concentration) can differentiate between cell growth and protein expression. It was demonstrated that the repression of the inducible invertase promoter in the presence of glucose and derepression by exhaustion of glucose could be utilized in expression of the green fluorescent protein as a heterologous protein.

## EXAMPLE 14

Construction of inducible lipocortin I expression vector pRI0LPI

[0067] PCR using plasmid pINV3000 (Example 1) carrying the *S. pombe* invertase gene as the template and oligo DNAs shown in SEQ ID NOS: 14 and 15 in the Sequence Listing as primers gave an amplification product containing the promoter region in the invertase gene and having restriction enzyme recognition sequences at both ends. After terminal double restriction digestion with *SpeI* and *EcoRI*, the amplification product was subjected to agarose gel electrophoresis. Purification of the band of about 3000 bp by the glass beads method using EASY-TRAP gave a fragment for use as a promoter insert.

[0068] The expression vector pTL2L (JP-A-7-163373) carrying a human lipocortin I gene was subjected to agarose gel electrophoresis after terminal double restriction digestion with *EcoRI* and *HindIII*. Purification of the band of about 1000 bp by the glass beads method using EASY-TRAP gave a fragment for use as the OPF insert.

[0069] The *S. pombe* multicloning vector pTL2M (JP-A-7-163373) was subjected to agarose gel electrophoresis after terminal double restriction digestion with *SpeI* and *HindIII*. Purification of the band of about 4500 bp by the glass beads method using EASY-TRAP gave a vector fragment.

[0070] The three fragments were ligated with a DNA ligation kit and transformed into *E. coli* strain DH5. *E. coli* colonies were screened for the inducible lipocortin I expression vector pRI0LPI through base sequencing and restriction mapping, and the vector was recovered by the alkali-SDS method on a preparatory scale.

## EXAMPLE 15

Preparation of fission yeast *Schizosaccharomyces pombe* transformant ASP139

[0071] A *S. pombe* wild-type strain ARC001 was transformed with the inducible lipocortin I expression vector pRI0LPI and a transducing vector pAL7 as described by Okazaki et al.

[0072] 1 ml of a preincubated ARC001 culture in YPD medium was incubated in minimum medium MB+Leu with shaking at 30°C for 16 hours to a cell density of  $1 \times 10^7$  per 1 ml, and the cells were collected, washed with water, suspended in 1 ml of 0.1M lithium acetate (pH 5.0) and incubated at 30°C for 60 minutes. A 100 µl portion of the suspension was mixed with 2 µg of the recombinant vector pRI0LPI and 0.5 µg of PstI-digested pAL7 in 15 µl TE and further with 290 µl of 50% PEG 4000 thoroughly and incubated at 30°C for 60 minutes, at 43°C for 15 minutes and at room temperature for 10 minutes, successively. After centrifugal removal of PEG, the cells were suspended in 1 ml of 1/2 YEL+Leu medium. After 10-fold dilution, 1 ml of the suspension was incubated at 32°C for 2 hours, and a 300 µl portion was spread on minimum medium agar MMA. After 3 days of incubation at 32 °C, about 300 independent colonies had developed on the plate.

[0073] 10 colonies of the transformant were inoculated in 2 ml of YEL medium containing 10 µg/ml antibiotic G418 (YEL10 medium) and incubated with shaking at 32°C. 2 days later, 2 clones were viable. All the subcultures of them were viable 3 days later. The putative desired transformant (ASP138 strain) was frozen in glycerol and stored for use in subsequent experiments.

## EXAMPLE 16

Analysis of lipocortin I expression

[0074] *S. pombe* transformant ASP139 (Example 15) was incubated in YPD medium containing 100 µg/ml G418 (glucose concentration 8%) at 32°C to the stationary phase and collected as a non-inducible cell culture. Separately, ASP139 was incubated in the same medium at first to the mid-growth phase, then after medium change, incubated in YPDG medium containing 100 µg/ml G418 (glucose concentration 0.1%, glycerol concentration 3%) to the stationary phase and collected as an inducible cell culture. Both cell cultures were washed, suspended in 50mM tris-HCl (pH 7.5) and disrupted with glass beads in a mini bead beater. After removal of the glass beads, the cell extracts were heated in the presence of SDS (1%) at 80°C for 15 minutes.

[0075] 50 µg protein from each extract was separated by SDS-polyacrylamide gel electrophoresis and stained with Coomassie Brilliant Blue. The extract from the inducible cell culture showed a major band of a molecular weight of about 45000, which is the same as the deduced molecular weight of the recombinant lipocortin I protein, but the extract from the non-inducible cell culture did not. More sensitive western analysis of band density showed that the band from the inducible cell culture extract was 10 times denser than the band from the non-inducible cell culture extract. The results show that while lipocortin I was not expressed in the cells incubated without medium change, the expression of lipocortin I in the cells incubated with the medium change was activated by the medium change probably because the depletion of glucose in the medium provoked derepression of the invertase promoter and thereby induced the protein expression. The high level expression induced by the medium change to a low-glucose expression medium from a high-glucose medium suggests that use of a growth medium (with a high glucose concentration) and an expression medium (with a low glucose concentration) can differentiate between cell growth and protein expression. It was demonstrated that the repression of the inducible invertase promoter in the presence of glucose and derepression by exhaustion of glucose could be utilized in expression of lipocortin I as a heterologous protein.

## EXAMPLE 17

Construction of expression vector pTL2INV1 carrying invertase gene

[0076] PCR using plasmid pINV3000 (Example 1) carrying the *S. pombe* invertase gene as the template and oligo DNAs shown in SEQ ID NOS: 16 and 17 in the Sequence Listing as primers gave an amplification product containing the ORF in the invertase gene and having restriction enzyme recognition sequences at both ends. After terminal double restriction digestion with AflIII (New England Biolab) and HindIII (Takara Shuzo Co., Ltd), the amplification product was subjected to agarose gel electrophoresis. Purification of the band of about 3000 bp by the glass beads method using EASY-TRAP gave a fragment for use as an insert.

[0077] The *S. pombe* multicloning vector pTL2M (JP-A-7-163373) was subjected to agarose gel electrophoresis after terminal double restriction digestion with SpeI and EcoI. Purification of the band of about 4500 bp by the glass beads method using EASY-TRAP gave a vector fragment.

[0078] The two fragments were ligated with a DNA ligation kit (Takara Shuzo Co., Ltd) and transformed into *E. coli* strain DH5 (Toyobo Co., Ltd.). *E. coli* colonies were screened for the invertase gene expression vector pRI0LPI through base sequencing and restriction mapping, and the vector was recovered multiplied by the alkali-SDS method on a preparatory scale.

#### EXAMPLE 18

##### Construction of secretory expression vector pSL2I06a'c1 using the signal sequence from the invertase gene

[0079] PCR using plasmid pINV3000 (Example 1) carrying the *S. pombe* invertase gene as the template and oligo DNAs shown in SEQ ID NOS: 18 and 19 in the Sequence Listing as primers gave an amplification product containing the ORF in the invertase gene and having restriction enzyme recognition sequences at both ends. After terminal double restriction digestion with *SpeI* (Takara Shuzo Co., Ltd.) and *EcoRI* (Takara Shuzo Co., Ltd.), the amplification product was subjected to agarose gel electrophoresis. Purification of the band of about 700 bp by the glass beads method using EASY-TRAP (Takara Shuzo Co., Ltd.) gave a fragment for use as a signal insert.

[0080] PCR using plasmid pSL2P06a'c1 (WO96/23890) containing human interleukin 6a'c1 variant cDNA as the template and oligo DNAs shown in SEQ ID NOS: 20 and 21 in the Sequence Listing as primers gave an amplification product containing the interleukin 6a'c1 variant ORF. After terminal double restriction digestion with *EcoRI* and *HindIII* (Takara Shuzo Co., Ltd), the amplification product was subjected to agarose gel electrophoresis. Purification of the band of about 600 bp by the glass beads method using EASY-TRAP (Takara Shuzo Co., Ltd.) gave a fragment for use as a gene insert.

[0081] The *S. pombe* multicloning vector pTL2M (JP-A-7-163373) was subjected to agarose gel electrophoresis after terminal double restriction digestion with *SpeI* and *HindIII*. Purification of the band of about 4500 bp by the glass beads method using EASY-TRAP gave a vector fragment.

[0082] The three fragments were ligated with a DNA ligation kit and transformed into the *E. coli* strain DH5 (Toyobo Co., Ltd.). *E. coli* colonies were screened for the IL-6a'c1 secretory expression vector pSL2I06a'c1 through base sequencing and restriction mapping, and the vector was recovered by the alkali-SDS method on a preparatory scale.

#### EXAMPLE 19

##### Preparation of *S. pombe* transformant ASP168

[0083] A leucine-requiring *S. pombe* strain ARC001 was transformed with the interleukin-6a'c1 variant secretory expression vector pSL2I06a'c1 (Example 18) and a transducing vector pAL7 as described by Okazaki et al.

[0084] 1 ml of a preincubated ARC001 culture in YPD medium was incubated in 100 minimum medium MB+Leu with shaking at 30°C for 16 hours. The cells were collected, washed with water, suspended in 0.1M lithium acetate (pH 5.0) at a cell density of  $10^9$  cells/ml and incubated at 30°C for 60 minutes. A 100 µl portion of the suspension was mixed with 2 µg of the recombinant vector pSL2I06a'c1 and 1.0 µg of *PstI*-digested pAL7 in 15 µl TE and further with 290 µl of 50% PEG 4000 thoroughly and incubated at 30°C for 60 minutes, at 43°C for 15 minutes and at room temperature for 10 minutes, successively. After centrifugal removal of PEG, the cells were suspended in 1 ml of 1/2 YEL+Leu medium. After 10-fold dilution, 1 ml of the suspension was incubated at 32°C for 2 hours, and a 300 µl portion was spread on minimum medium agar MMA. After 3 days of incubation at 32 °C, about 1000 independent colonies had developed on the plate.

[0085] The transformants (colonies) were inoculated in 2 ml of YEL medium containing 10 µg/ml antibiotic G418 (YEL10 medium) and incubated with shaking at 32°C for 5 days. The viable clones of the putative desired transformant, designated as strain ASP168, were frozen in glycerol and stored for use in subsequent experiments.

#### EXAMPLE 20

##### Analysis of expressed secretory interleukin-6a'c1 variant in culture medium

[0086] A fission yeast *Schizosaccharomyces pombe* transformant ASP168 (Example 19) was incubated in MA-Casamino acid medium (MA medium containing 2% Casamino acid and 3% glucose; for the composition of MA medium, refer to "Alfa et al., Experiments with Fission Yeast, Cold Spring Harbor Laboratory Press, 1993") containing 500 µg/ml G418 at 32°C for 2 days.

[0087] The cell culture was centrifuged, and the supernatant was concentrated 100-fold through a membrane filter (Amicon Co., Ltd.). Analysis of the concentrated sample by SDS-polyacrylamide electrophoresis followed by Coomassie Brilliant Blue staining gave the SDS-PAGE pattern shown in Figure 15. Lane 1 is the purified interleukin-6a'c1

## EP 1 223 219 A2

variant, lane 2 is the supernatant from the ASP168 cell culture, and lane 3 is the supernatant from a cell culture of the control strain ASP021 [transformant carrying a recombinant vector with no ORF prepared by recombination of pTL2M (JP-A-7-163373) by the method disclosed in JP-A-7-163373 without introduction of any gene to be expressed]. The band with a molecular weight of about 20000 in lane 3 seemed attributable to the interleukin-6a'c1 variant from the comparison of lanes 1 and 3.

**[0088]** Further analysis by western blotting using an anti-IL-6a'c1 gave the pattern shown in Figure 16. Lane 1 is the purified interleukin-6a'c1 variant, lane 2 is the supernatant from the ASP168 cell culture, and lane 3 is the supernatant from a cell culture of the control strain ASP021. The band with a molecular weight of about 20000 in lane 3 was identified as the interleukin-6a'c1 variant from the comparison of lanes 1 and 3.

## SEQUENCE LISTING

&lt;110&gt; Asahi Glass Co., Ltd.

&lt;120&gt; Inducible promoter gene and secretion signal gene functional in Schizosaccharomyces pombe, expression vector containing them and their use

&lt;130&gt; AB23(F9865)

&lt;150&gt; JP9/314608

&lt;151&gt; 1997-10-31

&lt;160&gt; 21

&lt;170&gt; MS-DOS

&lt;210&gt; 1

&lt;211&gt; 4748

&lt;212&gt; DNA

&lt;213&gt; Schizosaccharomyces pombe

&lt;400&gt; 1

```

ggatcctagt ccgcgaaatc gagatgcttt gaagattzaa attaaattta attttatgcg      60
agactggttt ctttatTTTT tgtatagtcg catgcaagcg aggttcgcat aatttggaaa      120
ataaaggtag tcaagaagac gttgaattaa ggctgcagtt tcaaagtact ctacaaacga      180
ttccttttta aaaaaaagat tcaaaaaaaa ggcaaagggg ttaagtaatg cttgttattt      240
caatttacct ccaaacagtt actaatgcaa ttgcaaaaaa aaaacctacc tattgaatca      300
aaatttctag cccatccatc gctcctcaag ataaaggaat cgatatTTTg agtttaazggg      360
agttgctgat agatttcaga attaaaaatt ttTggaaaag gatgtcgaga acaagaagat      420
acgtctagat tgctgatgat gcattctagc agacggaaat acaacgatat gtggacagca      480
cgacttttga tccgttcgga tcaaaaggaa gagaaatata catctttcaa gaagaatgca      540
ggaaaagcaa taaatgcccc ttTgattcct aaattatccc caaaaatgaa cattatgaga      600
tcttctTgtg ggagacagga aatttcgcaa ttccaaacga aaattcggct ctttttttta      660
ccccacagtt gcggggtaaa tgatgtaacg gacctTgggg gaaaggatga tgagttagtt      720
gggaagcgga aaaaatggaa aacggaagta agaatagaaa ccagtatggc tgagtgcatt      780
ggcgggaaaag attttacaga gatgacaaga atctatttat ctataaggaa aaacttttc      840
caaatttgtc taaaaacgca ttctcctcaa ttgcctctag gtagatgata taacgaattg      900
gaacgagaca tcgctaaccg gttttctTtg taaatgacat ttTgtagtgg gagtaagttt      960
gaatggaggg atagacagat gaatagtatg agatagaaga atagtatata taatgatata      1020
gatgaacaaa taaaaattga aagaaaaaag aaattgtTgg ctcattTggT tcatacacat      1080
gtTggTtcat acaactttta cccatcgtaa gtattataag taaaaaatag agtacgaaza      1140
gctataagta gtgaagcaaa aaaatagaaa aatagaaaaa aaaatatata taaaaaaata      1200
taataaaaaa aaaactcata agagacgtaa aacacaagaa ttgtctatca ttTgttcttt      1260

```

EP 1 223 219 A2

	aagaagcacc accatctctg aaaactcttc attctctatt agcaaggacc ctcttcattc	1320
5	cttctctctt agaatctctt tcattataac gaattggata atcgcgaat aagaacacat	1380
	ccccaaaata cgatatatcg atccatcttt tactttgcct agcttattgc tgtacaattc	1440
	catttaaata gtttctcttc aagaaagatc gicaatggag gcgacaatat accggaattc	1500
10	aagttgcgga cacagagctt gaaaagactg ctttttgtat tgttttcaag taaatgaaac	1560
	tgagttttga agtctcaaaa tacatcttat gtattgaaca ttagaagaac atataagata	1620
15	gatcttgaga gctcaattca tcgacattct agccatcata ctgcgatctt agacattgtc	1680
	agcacaacct tagatcgaaa atgaacacgt taccaaacgt tgtctaaaac ttgccgaatc	1740
	ttatctccgc attacttccg taatccttag tacatacgtt gcaatttcgg aagggtcatga	1800
20	tcgacttttt gtgtagctat aagtgcgca aatgagaaac atgacaagggt gcgatattta	1860
	gcaagatatatt atgcatttga tggagaaaagg aaatttcgga tgtatatata gtaccgttag	1920
	ctgcgctttt ttgggtcctc cataattttc aaactcactg ctttcgatca gatttaccgt	1980
25	ttttaaggtc ttatttgcct tgtgatctgt aggttggaaac atctatagtt cattttctaa	2040
	aagatccttt catcgtttca tcggatagta atcgttcaag aaaaaaaaaag aaaaaaagaa	2100
30	aaagaaaaag aaaagaaaaa taaaccgcta taattcatta cctatttgac tgaaggttct	2160
	tcactttgaa ttgttttgaa tcaaaataaa gaaattatta ttattatttt tttctctgc	2220
	ttttctttta tccattcgtc gaaactattt ttctgtgat aaaagcaatc attccttttt	2280
35	cctgcttctc ttgttatctg aattttaaac gacttttttt cctcgtccat tccctaattc	2340
	tttgcgacct tttctgattc tacccttgggt ttgtacttcc gttgtgtaat tgttgagaaa	2400
	gtgaactgat tatttaattg ttgtgaaaaa aattctaaaa ctattttgtt tttcttgatc	2460
40	attcatcctt tgctcgcttg cttgaatat attcagaaatc gtctcccttt caacggaata	2520
	tgataatttg ttgaatactc taaatcaatt aacacctatc aaaagctgaa acattaaatc	2580
45	tattctcacc aaaaaaaaaag actcaagctt cttcgttgtt ggccgggtctc ttttttgttt	2640
	tacgattgtt aaattttata ctcaaacctg ccaattctcc acttttgact atttattgat	2700
	agtccttatt taattttctg ttcaccgatt atcgtctttt ttgtaataaa tctttcttgg	2760
50	aaccaaccaa ttaatacgtt ataactgcta actttgaaga ttgtctaca atg ttt ttg	2818
		Met Phe Leu
		1
55	aaz tat att tta gct agt ggc att tgc ctc gtc tct ctc tta tca tct	2866
	Lys Tyr Ile Leu Ala Ser Gly Ile Cys Leu Val Ser Leu Leu Ser Ser	



EP 1 223 219 A2

	5	10	15	
5	aca aac gcg gct ccc cgt cac tta tat gta aaa cgt tat cct gtc att Thr Asn Ala Ala Pro Arg His Leu Tyr Val Lys Arg Tyr Pro Val Ile 20 25 30 35	2914		
10	tat aat gct tcc aac atc act gaa gtc agc aat tct acc acc gtt cct Tyr Asn Ala Ser Asn Ile Thr Glu Val Ser Asn Ser Thr Thr Val Pro 40 45 50	2962		
15	cct cct cca ttc gta aat aca acg gcc cct aat ggg act tgt ttg ggt Pro Pro Pro Phe Val Asn Thr Thr Ala Pro Asn Gly Thr Cys Leu Gly 55 60 65	3010		
20	aac tat aac gag tat ctt cct tca gga tat tac aat gct acc gat cgt Asn Tyr Asn Glu Tyr Leu Pro Ser Gly Tyr Tyr Asn Ala Thr Asp Arg 70 75 80	3058		
25	ccc aaa att cat ttt act cct tct tcc ggt ttc atg aat gat cca aac Pro Lys Ile His Phe Thr Pro Ser Ser Gly Phe Met Asn Asp Pro Asn 85 90 95	3106		
30	gga ttg gta tat act ggc ggc gtc tat cac atg ttc ttc caa tat tca Gly Leu Val Tyr Thr Gly Gly Val Tyr His Met Phe Phe Gln Tyr Ser 100 105 110 115	3154		
35	cca aaa act cta aca gcc ggc gaa gtt cat tgg ggt cac aca gtt tcc Pro Lys Thr Leu Thr Ala Gly Glu Val His Trp Gly His Thr Val Ser 120 125 130	3202		
40	aag gat tta atc cat tgg gag aat tat cct att gcc atc tac ccc gat Lys Asp Leu Ile His Trp Glu Asn Tyr Pro Ile Ala Ile Tyr Pro Asp 135 140 145	3250		
45	gaa cat gaa aac gga gtt ttg tcc ctc cca ttt agt ggc agt gca gtc Glu His Glu Asn Gly Val Leu Ser Leu Pro Phe Ser Gly Ser Ala Val 150 155 160	3298		
50	gtc gat gtt cat aat tct tcc ggt ctc ttt tcc aac gac acc att ccg Val Asp Val His Asn Ser Ser Gly Leu Phe Ser Asn Asp Thr Ile Pro 165 170 175	3346		
55	gaa gag cgc att gtt tta att tat acc gat cat tgg act ggt gtt gct Glu Glu Arg Ile Val Leu Ile Tyr Thr Asp His Trp Thr Gly Val Ala 180 185 190 195	3394		
	gag cgt cag gct att gcg tat acc act gat ggt gga tat act ttc aaa Glu Arg Gln Ala Ile Ala Tyr Thr Thr Asp Gly Gly Tyr Thr Phe Lys 200 205 210	3442		
	aaa tat tca gga aat ccc gtt ctt gac att aat tca ctt caa ttc cgc Lys Tyr Ser Gly Asn Pro Val Leu Asp Ile Asn Ser Leu Gln Phe Arg 215 220 225	3490		
	gac ccc aag gta ata tgg gat ttc gat gct aat cgt tgg gtg atg att Asp Pro Lys Val Ile Trp Asp Phe Asp Ala Asn Arg Trp Val Met Ile	3538		

## EP 1 223 219 A2

	230	235	240	
5	gta gct atg tct caa aat tat gga att gcc ttt tat tcc tcc tat gac Val Ala Met Ser Gln Asn Tyr Gly Ile Ala Phe Tyr Ser Ser Tyr Asp 245 250 255	3586		
10	ttg att cac tgg acc gag tta tct gtt ttc tcc act tct ggt tat ttg Leu Ile His Trp Thr Glu Leu Ser Val Phe Ser Thr Ser Gly Tyr Leu 260 265 270 275	3634		
15	ggg ttg caa tat gaa tgc cct gga atg gct cgt gtg ccc gtt gaa ggc Gly Leu Gln Tyr Glu Cys Pro Gly Met Ala Arg Val Pro Val Glu Gly 280 285 290	3682		
20	acc gat gaa tac aaa tgg gta ctc ttc at ctc catc aat cct ggc gct Thr Asp Glu Tyr Lys Trp Val Leu Phe Ile Ser Ile Asn Pro Gly Ala 295 300 305	3730		
25	cca ttg gga gga tcc gtt gtc caa tac ttt gtt ggc gat tgg aat ggt Pro Leu Gly Gly Ser Val Val Gln Tyr Phe Val Gly Asp Trp Asn Gly 310 315 320	3778		
30	aca aac ttc gtc ccc gat gat ggc caa act aga ttc gta gac ttg ggt Thr Asn Phe Val Pro Asp Asp Gly Gln Thr Arg Phe Val Asp Leu Gly 325 330 335	3826		
35	aag gac ttt tac gcc agc gct ttg tat cac tcg tct tcc gcc aat gcc Lys Asp Phe Tyr Ala Ser Ala Leu Tyr His Ser Ser Ser Ala Asn Ala 340 345 350 355	3874		
40	gat gtt att gga gtt gga tgg gct agc aac tgg caa tac acc aac caa Asp Val Ile Gly Val Gly Trp Ala Ser Asn Trp Gln Tyr Thr Asn Gln 360 365 370	3922		
45	gct cct act caa gtt ttc cgc agt gct atg aca gtt gca cga aaa ttc Ala Pro Thr Gln Val Phe Arg Ser Ala Met Thr Val Ala Arg Lys Phe 375 380 385	3970		
50	act ctt cgc gac gtt cct cag aac ccc atg acc aac ctt act tct ctc Thr Leu Arg Asp Val Pro Gln Asn Pro Met Thr Asn Leu Thr Ser Leu 390 395 400	4018		
55	att caa acc cca ttg aat gtt tct ctc tta cga gat gaa aca cta ttt Ile Gln Thr Pro Leu Asn Val Ser Leu Leu Arg Asp Glu Thr Leu Phe 405 410 415	4066		
60	acc gca ccc gtt atc aat agt tca agt agt ctt tcg ggc tct ccg att Thr Ala Pro Val Ile Asn Ser Ser Ser Ser Leu Ser Gly Ser Pro Ile 420 425 430 435	4114		
65	act ctt cca agc aat acc gca ttc gag ttc aat gtc aca ctc agt atc Thr Leu Pro Ser Asn Thr Ala Phe Glu Phe Asn Val Thr Leu Ser Ile 440 445 450	4162		
70	aat tac aca gaa ggc tgc aca aca gga tat tgt ctg ggg cgt att atc Asn Tyr Thr Glu Gly Cys Thr Thr Gly Tyr Cys Leu Gly Arg Ile Ile	4210		

EP 1 223 219 A2

	455	460	465	
5	att gat cct gat gat cca tac aga tta caa tcc atc tcc gtg gac gtt Ile Asp Ser Asp Asp Pro Tyr Arg Leu Gln Ser Ile Ser Val Asp Val 470 475 480	4258		
10	gat ttt gca gct agc act tta gtc att aat cgt gcc aaa gct cag atg Asp Phe Ala Ala Ser Thr Leu Val Ile Asn Arg Ala Lys Ala Gln Met 485 490 495	4306		
15	gga tgg ttt aat tca ctt ttc acg cct tct ttt gcc aac gat att tac Gly Trp Phe Asn Ser Leu Phe Thr Pro Ser Phe Ala Asn Asp Ile Tyr 500 505 510 515	4354		
20	att tat gga aac gta act ttg tat ggt att gtt gac aat gga ttg ctt Ile Tyr Gly Asn Val Thr Leu Tyr Gly Ile Val Asp Asn Gly Leu Leu 520 525 530	4402		
25	gaa ctg tat gtc aat aat ggc gaa aaa act tac act aat gac ttt ttc Glu Leu Tyr Val Asn Asn Gly Glu Lys Thr Tyr Thr Asn Asp Phe Phe 535 540 545	4450		
30	ttc ctt caa gga gca aca cct gga cag atc agc ttc gct gct ttc caa Phe Leu Gln Gly Ala Thr Pro Gly Gln Ile Ser Phe Ala Ala Phe Gln 550 555 560	4498		
35	ggc gtt tct ttc aat aat gtt acc gtt acg cca tta aag act atc tgg Gly Val Ser Phe Asn Asn Val Thr Val Thr Pro Leu Lys Thr Ile Trp 565 570 575	4546		
40	aat tgc taaatatattt gtttcaagtt aggaaagtat aataactttt gtccttgcac Asn Cys 580	4602		
45	attcaattgt aaagtttagt ttatcctttc atcgtaacca caattgtcac ctaaatctct aaaaatctct tcacttatct agttaatgtc gtaacaaaaa agtccagtag cttcgggaza tgatgcttgg aatcatacaa gtcgac	4662 4722 4748		
50	<210> 2 <211> 7286 <212> DNA <213> Artificial Sequence <400> 2 agcttgaaaa aacctccac acctccccct gaacctgaaa cataaaatga atgcaattgt tgttgttaac ttgtttattg cagcttataa tgggttacaaa taaagcaata gcatcacaaa tttcacaaat aaagcatttt ttctactgca ttctagttgt gggttgtcca aactcatcaa tgtatcttat catgtctgga tcgatcccg gagggtgggc gtcgcttggc cgggtcatttc gaaccccaga gicccgctca gaagaactcg tcaagaaggc gatagaaggc gatgcgctgc	60 120 180 240 300		
55				

EP 1 223 219 A2

gaatcggag cggcgatacc gtaaagcācāg āggaagcgg cagccccattc gccgccaaagc 360  
 5 tcttcagcaa tatcacgggt agccaacgct atgtcctgat agcgggccgc cacacccagc 420  
 cggccacagt cgatgaatcc agaaaagcgg ccatcttcca ccatgatatt cggcaagcag 480  
 gcatcgccat gggtcacgac gagatcctcg ccgtcgggca tgcgcgcctt gagcctggcg 540  
 10 aacagttcgg ctggcgcgag cccctgatgc tcttcgtcca gatcatcctg atcgacaaga 600  
 ccggcttcca tccgagtagc tgcctcgctc atgcgatgtt tcgcttggtg gtcgaatggg 660  
 caggtagccg gatcaagcgt atgcagccgc cgcattgcat cagccatgat ggatactttc 720  
 15 tcggcaggag caaggtgaga tgacaggaga tcttgccccg gcacttcgcc caatagcagc 780  
 cagtccttcc cgccttcagt gacaacgtcg agcacagctg cgcaagggaac gcccgtcgtg 840  
 20 gccagccacg atagccgcgc tgcctcgctc tgcagttcat tcagggcacc ggacaggtag 900  
 gtcttgacaa aaagaaccgg gcgcccctgc gctgacagcc ggaacacggc ggcatcagag 960  
 cagccgattg tctgttggtc ccagtcatag ccgaatagcc tctccacca agcggccgga 1020  
 25 gaacctgcgt gcaatccatc ttgttcaatc atgcgaaacg atcctcatcc tgtctcttga 1080  
 tcagatccgg gacctgaaat aaaagacaaa aagactaaac ttaccagtta actttctggt 1140  
 ttttcagttc ctcgaggagc tttttgcaaa agcctaggcc tccaaaaaag cctcctcact 1200  
 30 acttctggaa tagctcagag gccgaggcgg cctcggcctc tgcataaata aaaaaatta 1260  
 gtcagccatg gggcgagaa tgggcggaac tgggcggagt tagggcggg atggcgggag 1320  
 35 ttagggcggg gactatggtt gctgactaat tgagatgcat gctttgcata cttctgcctg 1380  
 ctggggagcc tggggacttt ccacacctgg ttgctgacta attgagatgc atgctttgca 1440  
 tacttctgcc tgcctggggag cctggggact tccacaccc taactgacac acattccaca 1500  
 40 ggacattgat tattgactag ttagtccgcg aaatcgagat gctttgaaga ttaaaattaa 1560  
 atttaatttt atgcgagact ggtttcctta ttttttgat agtcgcatgc aagcgaggtt 1620  
 cgcataattt ggaaaaataa ggtagtcaag aagacgttga attaaggctg cagtttcaaa 1680  
 45 gtactctaca aacgattcct tttaaaaaaa aagattcaaa aaaaaggcaa agggtttaag 1740  
 taatgcttgt tatttcaatt tacctccaaa cagttactaa tgcaattgca aaaaaaaaaac 1800  
 50 ctacctattg aatcaaaatt tctagcccat ccacgctcc tcaagataaa ggaatcgata 1860  
 ttttgagttt aaggagttg ctgatagatt tcagaattaa aaatttttgg aaaaggatgt 1920  
 cgagaacag aagatacgtc tagattgctg atgatgcatt ctagcagacg gaaatacaac 1980  
 55

EP 1 223 219 A2

5      gataatgtgga cagcacgact ttgatccgt tcgzaatcaa aggaagagaa atatccatct 2040  
 ttcaagaaga atgcaggaaa agcaataaa: gccatttga ttcttaaat atccccaaaa 2100  
 10      atgaacatta tgagatcttc ttgtgggaga caggzaattt cgcaattcca aacgazaatt 2160  
 cggcictttt ttttacccca cagttgcggg gtaaatgatg taacggacct tgggggzaag 2220  
 gatgatgagt tagttgggaa gcggaaaaaa tggaaaacgg aagtaagaat agaaaccagt 2280  
 atggctgagt gcaatggcgg aaaagatttt acagagatga caagaatcta tttatctata 2340  
 15      aggaaaaact ttttccaaat ttgtctaaaa acgcattctc ctcaattgcc tctaggtaga 2400  
 tgatataacg aattggaacg agacatcgct aaccggtttt ctttgtaaat gacattttgt 2460  
 agtggggagta agtttgaatg gagggataga cagatgaata gtatgagata gaagaatagt 2520  
 atatataatg attaagatga acaataaaaa attgaaagaa aaaagaaatt gttggctcat 2580  
 20      ttggttcata cacatgttgg ttcatataac ttttaccat cgtaagtatt ataagtaaaa 2640  
 aatagagtac gaaaagctat aagtagtgaa gcaaaaaaat agaaaaatag aaaaaaaat 2700  
 25      atatataaaa aaatataata aaaataaaac tcataagaga cgtaaaacac aagaattgtc 2760  
 tatcatttgt tctttaagaa gcaccacat tctgtaaaac tcttcatttc tcattagcaa 2820  
 ggaccctttt cattccttcc tctttagaat ccttttcatt ataacgaatt ggataatacg 2880  
 30      caaataagaa cacatccctt aaatacgata tatcgatcca ttttttactt tgcctagctt 2940  
 attgctgtac aattccattt aaatagtttc tcttcaagaa agatcgtaa tggaggcgac 3000  
 aatataccgg aatttaagtt gcggacacag agcttgaaaa gactgcattt tgtattgttt 3060  
 35      tcaagtaaat gaaactgagt ttggaagtct caaaatacat cttatgtatt gaacattaga 3120  
 agaacatata agatagatct tgagagctca attcatcgac attctagcca tcatactgcg 3180  
 40      atcttagaca ttgtcagcac aaccttagat cgaaaatgaa cagttacca aacgttgtct 3240  
 aaaacttgcc gaattctatc tccgcattac tccgtaatc cttagtacat acgctgcaat 3300  
 ttcggaaggt catgatcgac tttttgtgta gctataagtg acgcaaatga gaaacatgac 3360  
 45      aaggtgcgat atttagcaag atattatgca ttgatggag aaaggaaatt tcggatgtat 3420  
 atatag:acc gttagctgcg ctttttttgg tcatacataa ttttcaaact cactgcttcc 3480  
 50      gatcagattt accgttttta aggtctttat tgccttgtga tctgtaggtt ggaacatcta 3540  
 tagttcattt tctaaaagat cctttcatcg ttcatcgga tagtaatcgt tcaagaaaaa 3600  
 aaaagaaaaa aagaaaaaga aaaagaaaag aaaaataaac cgctataatt cattacctat 3660  
 55

EP 1 223 219 A2

	ttgactgaag gtctctcatc ttgaattgtt ttgaatcaaa ataaagaaat tarrattatt	3720
5	tatttttttc ttgccttttc ctttatccat tgcgcgaaac tattttctcg ctgataaaag	3730
	caatcattcc tttttcctgc ttctcttgtt attcgaattt taaacgactt tttttcctcg	3840
	tccattccct aattctttgc gaccttttct gattctatcc ttggtttgta ctttcgttgt	3900
10	gtaattgttg agaaagtga ctagattattt aattgtttgtg aaaaaaatc taaaactatt	3960
	ttgtttttct tgatcattca tcccttgctc gcttgcttga atattacaga aattcgtctc	4020
15	cctttcaacg gaatatgata atttgttgaa tactctaaat caattaacac ctatcaaaag	4080
	ctgaaacatt aaatctattc tcaccaaaaa aaaagactca agcttcttcg ttgttgcccg	4140
	gtctcttttt tgttttacga ttgttaaat ttatactcac aactgccaat tctccacttt	4200
20	tgactattta ttgatagtc ctatttaatt ttctgttcac cgattatcgt cttttttgta	4260
	aataatcttt cttggaacca accaattaat acgttataat cgctaacttt gaagatttgc	4320
25	tacaatggca atgggtacag aattcgagct cggtagccgg ggatcctcta gagtcgacct	4380
	gcaggcatgc aagcttaaat aggaaagttt ctcaacagg attacagtgt agctacctac	4440
	atgctgaaaa atatagcctt taaatcattt ttatattata actctgtata atagagataa	4500
30	gtccattttt taaaaatgtt tccccaaac cataaaacc tatacaagtt gttctagtaa	4560
	caatacatga gaaagatgtc tatgtagctg aaaataaaat gacgtcacaa gacaaaaaaa	4620
	aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaagta	4680
35	ccttctgagg cggaaagaac cagccggatc cagacatgat aagatacatt gatgagtttg	4740
	gacaaaccac aactagaatg cagtgaaaaa aatgctttat ttgtgaaatt tgtgatgcta	4800
	ttgctttatt tgaaccatt ataagctgca ataaacaagt taacaacaac aattgcattc	4860
40	attttatgtt tcagggtcag ggggaggtgt gggaggtttt ttaaagcaag taaaacctct	4920
	acaaatgttg tatggctgat tatgatccgg ctgcctcgcg cgtttcggtg atgacgggtga	4980
45	aaacctctga cacatgcagc tcccgagagc ggtagacagt tgtctgtaag cggatgccgg	5040
	gagcagacaa gcccgtcagg gcgcgtcagc ggggtgttggc ggggtgtcggg gcgcagccat	5100
	gacccagtc cgtagcgata gcggagtgt tactggctta actatgcggc atcagagcag	5160
50	atigtactga gagtgcacca tatgcggtgt gaaataccgc acagatgcgt aaggagaaaa	5220
	taccgcatca ggcgtcttc cgttctctcg ctactgact cgctgcgctc ggtcgttcgg	5280
55	ctgcggcgag cggtatcagc tcactcaaa ggcgtaatac ggttatccac agaatacagg	5340

EP 1 223 219 A2

gataacgcag gaaagaatga gcaaaaggcc agcaaaaggc caggaaccgt aaaaaggccg 5400  
 5 cgttgcctggc gtttttccat aggtcccgcc cccctgacga gcatcacaaa aatcgacgct 5460  
 caagtcagag gtggcgaaac ccgacaggac tataaagata ccaggcggtt ccccttgga 5520  
 gctccctcgt gcgctctcct gtcccgaccc tgccgcttac cggataacctg tccgcctttc 5580  
 10 tcccttcggg aagcgtggcg ctttctcata gctcacgctg taggtatctc agttcgggtg 5640  
 aggtcgttcg ctccaagctg ggctgtgtgc acgaaccccc cgttcagccc gaccgctgcg 5700  
 ccttatccgg taactatcgt cttgagtcga acccggttaag acacgactta tcgccactgg 5760  
 15 cagcagccac tggtaacagg attagcagag cgaggatatgt aggcgggtgct acagagtctt 5820  
 tgaagtgggtg gcctaactac ggctacacta gaaggacagt atttgggtatc tgcgctctgc 5880  
 20 tgaagccagt taccttcgga aaaagagttg gtagctcttg atccggcaaa caaaccaccg 5940  
 ctggtagcgg tgggtttttt gtttgcaagc agcagattac gcgcagaaaa aaaggatctc 6000  
 aagaagatcc tttgatcttt tctacggggt ctgacgctca gtggaacgaa aactcacgtt 6060  
 25 aagggatttt ggtcatgaga ttatcaaaaa ggatcttcac ctagatcctt ttaaattaaa 6120  
 aatgaagttt taaatcaatc taaagtatat atgagtaaac ttggctctgac agttaccaat 6180  
 gcttaatcag tgaggcacct atctcagcga tctgtctatt tcgttcatcc atagtgcct 6240  
 30 gactccccgt cgtgtagata actacgatac gggaggggctt accatctggc cccagtgcgtg 6300  
 caatgatacc gcgagacca cgctcaccgg ctccagattt atcagcaata aaccagccag 6360  
 35 ccggaagggc cgagcgcaga agtggctctg caactttatc cgcctccatc cagtctatta 6420  
 attgttgccg ggaagctaga gtaagtagtt cgccagttaa tagtttgccg aacgttggtg 6480  
 ccattgtctg aggcategtg gtgtcacgtc cgtcgtttgg tatggcttca ttcagctccg 6540  
 40 gttcccaacg atcaaggcga gttacatgat ccccatgtt gtgcaaaaaa gcggttagct 6600  
 ccttcggtcc tccgatcgtt gtcagaagta agttggccgc agtggttatca ctcatggtta 6660  
 tggcagcact gcataattct cttactgtca tgccatccgt aagatgcttt tctgtgactg 6720  
 45 gtgagtactc aaccaagtca ttctgagaat agtgtatgcg gcgaccgagt tgctcttgcc 6780  
 cggcgtcaac acgggataat accgcgccac atagcagaac ttlaaaagtg ctcatcattg 6840  
 50 gaaaacgttc ttcggggcga aaactctcaa ggatcttacc gctgttgaga tccagttcga 6900  
 tgtaaccac tcgtgcaccc aactgatctt cagcatcttt tacittcacc agcgtttctg 6960  
 ggtgagcaaa aacaggaagg caaaatgccg caaaaaaggg aataagggcg acacggaaat 7020

55



EP 1 223 219 A2

gttgaatact cataactcttc ctttttcaat attattgaag catttatcag ggttatgtc 7080  
5 tcatgagcgg atacatatct gaatgtatct agaaaaataa acaaataagg gttccgcgca 7140  
catttccccg aaaagtgcga cctgacgtct aagaaacat tattatcatg acattaacct 7200  
ataaaaaatag gcgtatcacg aggcccttc gtcttcaaga attggtcgac caattctcat 7260  
10 gtttgacagc ttatcatcga taagct 7286

<210> 3  
15 <211> 27  
<212> DNA  
<213> Artificial Sequence  
<400> 3  
gtttgaattc atgaaygayg gnaaygg 27

20 <210> 4  
<211> 24  
<212> DNA  
<213> Artificial Sequence  
25 <400> 4  
gtttgaattc yrytggggnc aygc 24

30 <210> 5  
<211> 27  
<212> DNA  
<213> Artificial Sequence  
<400> 5  
gtttgaattc acyttnnggrt cncgraa 27

35 <210> 6  
<211> 24  
<212> DNA  
<213> Artificial Sequence  
40 <400> 6  
gatgggtaaa actagtatga acca 24

45 <210> 7  
<211> 23  
<212> DNA  
<213> Artificial Sequence  
<400> 7  
ttgcgaatta ctagtctccc aca 23

50 <210> 8  
<211> 29  
<212> DNA  
<213> Artificial Sequence  
55 <400> 8

# EP 1 223 219 A2

	gggactagtt agtccgcgaa atcgagatg	29
5	<210> 9 <211> 42 <212> DNA <213> Artificial Sequence <400> 9	
10	gaggaattct gataccattg ccattgtagc aaatcttcaa ag	42
15	<210> 10 <211> 30 <212> DNA <213> Artificial Sequence <400> 10	
	aaagctagcc attgtagcaa atcttcaaag	30
20	<210> 11 <211> 27 <212> DNA <213> Artificial Sequence <400> 11	
25	atggctagca agggcgagga gctgttc	27
30	<210> 12 <211> 27 <212> DNA <213> Artificial Sequence <400> 12	
	cccaagctta cttgtacagc tcgtcca	27
35	<210> 13 <211> 7938 <212> DNA <213> Artificial Sequence <400> 13	
40	agcttgaaaa aacctccac acctccccc gaacctgaaa cataaaatga atgcaattgt	60
	tgttgttaac ttgtttattg cagcttataa tgggtacaaa taaagcaata gcatcacaaa	120
	tttcacaaat aaagcatttt tttcactgca ttctagtgtt gggttgtcca aactcatcaa	180
45	tgtatcttat catgtctgga tcatcccg gagggtgggc gtcgcttggc cggctatttc	240
	gaacccaga gtcccgctca gaagaactcg tcaagaaggc gatagaaggc gatgcgctgc	300
50	gaatcggggag cggcgatacc gtaaagcacg aggaagcggc cagcccatte gccgccaagc	360
	tcttcagcaa tatcaggggt agccaacgct atgtcctgat agcggtcgc caccaccagc	420
	cggccacagt ccatgaatcc agaaaagcgg ccattttcca ccatgatatt cggcaagcag	480
55		

## EP 1 223 219 A2

5  
 10  
 15  
 20  
 25  
 30  
 35  
 40  
 45  
 50  
 55

gcatcgccat gggtcacgac gagatccctcg ccgtcgggca tgcgcgcctt gagcctggcg 540  
 aacagttcgg ctggcgcgag cccctgatgc tcttcgtcca gatcatcctg atcgacaaga 600  
 ccggcttcca tccgagtagc tgcctcgctcg atgcgatgtt tcgcttgggtg gtcgaatggg 660  
 caggtagccg gatcaagcgt atgcagccgc cgcattgcat cagccatgat ggatactttc 720  
 tcggcaggag caaggtgaga tgacaggaga tccctgccccg gcacttcgcc caatagcagc 780  
 cagtccttcc ccgcttcagt gacaacgtcg agcacagctg cgcaagggaac gcccgtcgtg 840  
 gccagccacg atagccgcgc tgcctcgctc tgcagttcat tcagggcacc ggacaggctg 900  
 gtcttgacaa aaagaaccgg gcgccccctgc gctgacagcc ggaacacggc ggcatcagag 960  
 cagccgattg tctgtttgtc ccagtcatag ccgaatagcc tctccacca agcggccgga 1020  
 gaacctgcgt gcaatccatc ttgttcaatc atgcgaaacg atcctcatcc tgtctcttga 1080  
 tcagatccgg gacctgaaat aaaagacaaa aagactaaac ttaccagtta actttctggg 1140  
 ttttcagttc ctcgaggagc tttttgcaaa agcctaggcc tccaaaaaag cctctcact 1200  
 actttctgaa tagctcagag gccgaggcgg cctcggcctc tgcataaata aaaaaaatta 1260  
 gtcagccatg gggcggagaa tgggcggaac tgggcggagt tagggcgggg atgggcggag 1320  
 ttaggggcgg gactatgggt gctgactaat tgagatgcat gctttgcata cttctgcctg 1380  
 ctggggagcc tggggacttt ccacacctgg ttgctgacta attgagatgc atgctttgca 1440  
 tacttctgcc tgcitggggag cctggggact ttccacaccc taactgacac acattccaca 1500  
 ggacattgat tattgactag ttagtccgcg aaatcgagal gctttgaaga ttaaaattaa 1560  
 atttaatttt atgcgagact ggtttcctta tttttgtat agtcgcatgc aagcgaggtt 1620  
 cgcataattt ggaaaataaa ggtagtcaag aagacgttga attaaggctg cagtttcaaa 1680  
 gtactctaca aacgattcct tttaaaaaaa aagattcaaa aaaaaggcaa agggtttaag 1740  
 taatgcttgt tattttcaatt tacctccaaa cagttactaa tgcaattgca aaaaaaaaaac 1800  
 ctacctattg aatcaaaatt tctagcccat ccacgcctcc tcaagataaa ggaatcgata 1860  
 ttttgagttt aaggaggttg ctgatagatt tcagaattaa aaatttttgg aaaaggatgt 1920  
 cgagaacaag aagatacgtc tagattgctg atgatgcatt ctacgagacg gaaatacaac 1980  
 gatatttgga cagcacgact ttgatccgt tcggatcaaa aggaagagaa atatccatct 2040  
 ttcaagaaga atgcaggaaa agcaataaat gcccatttga ttcctaaatt atccccaaaa 2100  
 atgaacatta tgagatcttc ttgtgggaga caggaaattt cgcaattcca aacgaaaatt 2160

EP 1 223 219 A2

5  
 10  
 15  
 20  
 25  
 30  
 35  
 40  
 45  
 50  
 55

cggcctctttt ttttacccca cagttgcggg gtaaatgatg taacggacct tgggggaaaag 2220  
 gatgatgagt tagttgggaa gcggaaaaaa tggaaaacgg aagtaagaat agaaaccagt 2280  
 atggctgagt gcaatggcgg aaaagatttt acagagatga caagaatcta tttatctata 2340  
 aggaaaaact ttttccaaat ttgtctaaaa acgcattctc ctcaattgcc tctaggtaga 2400  
 tgatataacg aattggaacg agacatcgct aaccggtttt ctttgtaaat gacattttgt 2460  
 agtgggagta agtttgaatg gagggataga cagatgaata gtatgagata gaagaatagt 2520  
 atatataatg attaagatga acaataaaaa attgaaagaa aaaagaaatt gttggctcat 2580  
 ttggttcata cacatgttgg ttcatacaac ttttaccat cgtaagtatt ataagtaaaa 2640  
 aatagagtac gaaaagctat aagtagtgaa gcaaaaaaat agaaaaatag aaaaaaaat 2700  
 atatataaaa aaatataata aaaataaaac tcataagaga cgtaaaacac aagaattgtc 2760  
 tatcatttgt tctttaagaa gcaccaccat tctgtaaaac tcttcatttc tcattagcaa 2820  
 ggaccctttt cattccttcc tctttagaat ctttttcatt ataacgaatt ggataatacg 2880  
 caaataagaa cacatccctt aaatacgata tatcgatcca ttttttactt tgcctagctt 2940  
 attgctgtac aattccattt aaatagtttc tcttcaagaa agatcgtaa tggaggcgac 3000  
 aatataccgg aatttaagtt gcggacacag agcttgaaaa gactgcattt tgtattgttt 3060  
 tcaagtaaat gaaactgagt ttggaagctt caaaatacat cttatgtatt gaacattaga 3120  
 agaacatata agatagatct tgagagctca attcatcgac attctagcca tcatactgcg 3180  
 atcttagaca ttgtcagcac aaccttagat cgaaaatgaa cacgttacca aacgttgtct 3240  
 aaaacttgcc gaatcttate tccgcaftac tccgtaatc cttagtacat acgctgcaat 3300  
 ttcggaaggt catgatcgac tttttgtgta gctataagtg acgcaaatga gaaacatgac 3360  
 aagggtcgat atttagcaag atattatgca ttgatggag aaaggaaatt tcggatgcat 3420  
 atatagtacc gtttagctcg ctttttttgg tcatccataa ttttcaaact cactgctttc 3480  
 gatcagattt accgttttta aggtctttat tgctttgtga tctgtagggt ggaacatcta 3540  
 tagttcattt tctaaaagat cttttcctcg tttcatcgga tagtaatcgt tcaagaaaaa 3600  
 aaaagaaaaa aagaaaaaga aaaagaaaag aaaaataaac cgctataatt cattacctat 3660  
 ttgactgaag gttcttcac ttgaattgtt tgaatcaaa ataaagaaat tattattatt 3720  
 attttttttc ttcgcttttt ctttatccat tcgtcgaaac tatttttctg ctgataaaaag 3780  
 caatcattcc tttttctgc ttctcttgtt attcgaaatt taaacgactt tttttctcg 3840

EP 1 223 219 A2

tccattccct aattctttgc gaccttttct gattctatcc ttggtttga ctttcgttgc 3900  
 gtaattgttg agaaagtga ctagttattt aattgttttg aaaaaaattc taaaactatt 3960  
 5 ttgtttttct tgatcatcca tcttttgctc gcttgcttga atattacaga aattcgtctc 4020  
 cctttcaacg gaatatgata atttgttgaa tactctaaat caattaacac ctatcaaaag 4080  
 10 ctgaaacatt aaatctattc tcaccaaaaa aaaagactca agcttcttcg ttgttggccg 4140  
 gtctcttttt tgttttacga ttgttaaatt ttatactcac aactgccaat tctccacttt 4200  
 tgactattta ttgatagtcc ctatttaatt tctgtttcac cgattatcgt ctttttttga 4260  
 15 aataatcttt ctltggaacca accaattaat acgttataat cgctaacttt gaagatttgc 4320  
 tacaatggct agcaagggcg aggagctgtt caccgggggtg gtgcccaccc ttggtcgagct 4380  
 ggacggcgac gtaaacggcc acaagttcag cgtgtccggc gagggcgagg gcgatgccac 4440  
 20 ctacggcaag ctgacctga agttcatctg caccaccggc aagctgcccg tgccttgccc 4500  
 caccctcgtg accaccctga cctacggcgt gcagtgtctc agccgctacc ccgaccacat 4560  
 25 gaagcagcac gactttctca agtccgccat gcccgaggc tacgtccagg agcgcaccat 4620  
 cttcttcaag gacgacggca actacaagac ccgcgccgag gtgaagtctg agggcgacac 4680  
 cctggtgaac cgcatcgagc tgaagggcat cgacttcaag gaggacggca acatcctggg 4740  
 30 gcacaagctg gagtacaact acaacagcca caacgtctat atcatggccg acaagcagaa 4800  
 gaacggcatc aagggtgaact tcaagatccg ccacaacatc gaggacggca gcgtgcagct 4860  
 35 cgccgaccac taccagcaga acacccccat cggcgacggc cccgtgctgc tgcgcgacaa 4920  
 ccactacctg agcaccagct ccgccctgag caaagacccc aacgagaagc gcgatcacat 4980  
 ggtcctgctg gagttcgtga ccgccgccgg gatcactctc ggcatggacg agctgtacaa 5040  
 40 gtaagcttaa ataggaaggt tctttcaaca ggattacagt gtagctacct acatgctgaa 5100  
 aaatatagcc tttaaatcat ttttatatta taactctgta taatagagat aagtccattt 5160  
 tttaaaaatg ttttcccaa accataaaac cctatacaag ttgttctagt aacaatacat 5220  
 45 gagaagatg tctatgtagc tgaaaataaa atgacgtcac aagacaaaaa aaaaaaaaaa 5280  
 aaaaaaaaaa aaaaaaaaaa aaaaaaaa aaaaaaaaaa aaaaaaaaag taccttctga 5340  
 50 ggcggaaga accagccgga tccagacatg ataagatata ttgatgagtt tggacaaacc 5400  
 acaactagaa tgcagtga aaatgtctt atttgtgaaa ttgtgtatgc tattgcttta 5460  
 ttgttaacca ttataagctg caataaaca gttacaaca acaattgcat tcattttatg 5520

55

EP 1 223 219 A2

5 5580 tttcagggttc agggggagggt gtgggagggtt ttttaaagca agtaaaaccc ctacaaatgt  
 5640 ggtagggctg attatgatcc ggctgcctcg cgcgtttcgg tgatgacggg gaaaacctct  
 5700 gacacatgca gctcccggag acggtcacag cttgtctgta agcggatgcc gggagcagac  
 5760 aagcccgtca gggcgcgtca gcgggtgttg gcgggtgtcg gggcgcagcc atgaccagt  
 10 5820 cacgtagcga tagcggagtg tatactggct taactatgcg gcatcagagc agattgtact  
 5880 gagagtgcac catatgcggg gtgaaatacc gcacagatgc gtaaggagaa aataccgcat  
 5940 caggcgtctt tccgtttcct cgctcactga ctgcctgcgc tcggctcgtt ggctgcggcg  
 15 6000 agcggtatca gctcactcaa aggcggtaat acggttatcc acagaatcag gggataacgc  
 6060 aggaagaat gagcaaaagg ccagcaaaag gccaggaacc gtaaaaaggc cgcgttgcgt  
 20 6120 gcgtttttcc ataggctccg ccccccgtac gagcatcaca aaaatcgacg ctcaagtcag  
 6180 aggtggcgaa acccgacagg actataaaga taccaggcgt tccccctgg aagctccctc  
 6240 gtgcgtcttc ctgttccgac cctgcgcgtt accggatacc tgtccgcctt tctcccttcg  
 25 6300 ggaagcgtgg cgttttctca tagctcacgc ttaggtatc tcagttcggg ttaggtcgtt  
 6360 cgctccaagc tgggctgtgt gcacgaacc cccgttcagc ccgaccgtg cgccttatcc  
 6420 ggtaactatc gtcttgagtc caaccggta agacacgact tatcgccact ggcagcagcc  
 30 6480 actggtaaca ggattagcag agcggaggtat gtaggcggtg ctacagagtt cttgaagtgg  
 6540 tggcctaact acggctacac tagaaggaca gtttttgta tctgcgtctt gctgaagcca  
 35 6600 gttaccttcg gaaaaagagt tggtagctct tgatccggca aacaaaccac cgctggtagc  
 6660 ggtgggtttt ttgtttgcaa gcagcagatt acgcgcagaa aaaaaggatc tcaagaagat  
 6720 cctttgatct tttctacggg gtctgacgct cagtggaaag aaaactcacg ttaagggatt  
 40 6780 ttggatcatga gattatcaaa aaggatcttc acctagatcc ttttaaatta aaaatgaagt  
 6840 tttaaatcaa tctaaagtat atatgagtaa acttgggtctg acagttacca atgcttaatc  
 45 6900 agtgaggcac ctatctcagc gatctgtcta tttcggtcat ccatagttgc ctgactcccc  
 6960 gtcgttaga taactacgat acgggagggc ttaccatctg gccccagtgc tgcaatgata  
 7020 ccgcgagacc cacgctcacc ggctccagat ttatcagcaa taaaccagcc agccggaagg  
 50 7080 gccgagcgca gaagtgggtc tgcaacttta tccgcctcca tccagttctat taattgttgc  
 7140 cgggaagcta gagtaagtag ttccgaggtt aatagtttgc gcaacgttgt tgccattgct  
 7200 gcaggcatcg tgggtgtcag ctgcgtcgtt ggtatggctt cattcagctc cggttcccaa  
 55

EP 1 223 219 A2

5  
 10  
 15  
 20  
 25  
 30  
 35  
 40  
 45  
 50  
 55

cgaacaaaggc gagttacatg atcccccatg ttgtgcaaaa aagcggtag ctccctcggg 7260  
 cctccgatcg ttgtcagaag taagtggcc gcagtgttat cactcatggg tatggcagca 7320  
 ctgcataatt ctcttactgt catgccatcc gtaagatgct tttctgtgac tgggtgagtac 7380  
 tcaaccaagt cattctgaga atagtgtatg cggcgaccga gttgctcttg cccggcgctca 7440  
 acacgggata ataccgcgcc acatagcaga actttaaaag tgctcatcat tggaaaacgt 7500  
 tcttcggggc gaaaactctc aaggatctta ccgctgttga gatccagttc gatgtzaacc 7560  
 actcgtgcac ccaactgac ttcagcatct tttactttca ccagcgttc tgggtgagca 7620  
 aaaacaggaa ggcaaaatgc cgcaaaaaag ggaataaggc cgacacggaa atgttgaata 7680  
 ctcatactct tcctttttca atattattga agcatttctc agggttattg tctcatgagc 7740  
 ggatacatat ttgaatgtat ttagaaaaat aaacaaatag gggttccgcg cacatttccc 7800  
 cgaaaagtgc cacctgacgt ctaagaaacc attattatca tgacattaac ctataaaaaat 7860  
 aggcgtatca cgaggccctt tcgtcttcaa gaattggtcg accaattctc atgtttgaca 7920  
 gcttatcacc gataagct 7938  
 <210> 14  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence  
 <400> 14  
 gggactagtt agtccgcgaa atcgagatg 29  
 <210> 15  
 <211> 42  
 <212> DNA  
 <213> Artificial Sequence  
 <400> 15  
 gaggaattct gataccattg ccattgtagc aaatcttcaa ag 42  
 <210> 16  
 <211> 28  
 <212> DNA  
 <213> Artificial Sequence  
 <400> 16  
 gggac atgtt tttga aatat atttt agc 28  
 <210> 17  
 <211> 27  
 <212> DNA  
 <213> Artificial Sequence  
 <400> 17



gggaa gctta gcaat tccag atagt ct

27

5

<210> 18  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <400> 18  
 ttgac tagtt attaa tagta

20

10

15

<210> 19  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence  
 <400> 19  
 cccga attca taacg tttta catat aagt

29

20

25

<210> 20  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence  
 <400> 20  
 aaaga attcc cagta ccccc aacct cttca

29

30

35

<210> 21  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <400> 21  
 aaaat gattt aaagg ctata

20

40

#### Claims

1. A peptide having the amino acid sequence of Met Phe Leu Lys Tyr Ile Leu Ala Ser Gly Ile Cys Leu Val Ser Leu Leu Ser Ser Thr Asn Ala.

45

2. A DNA encoding the peptide of claim 1.

3. A recombinant vector containing the sequence of the DNA according to claim 2.

50

4. A multicloning vector containing the sequence of the DNA according to claim 2 and a multicloning site.

5. An expression vector containing the sequence of the DNA according to claim 2 and a heterologous protein structural gene.

55

6. A *Schizosaccharomyces pombe* transformant carrying the expression vector according to claim 5.

7. A process for producing a protein which comprises incubating the transformant according to claim 6 and recovering an expressed heterologous protein.

## F I G . 1

```

S. pombe inv1      31 : TTAPNGTCLGNYNEYLPSCGYINATDRPKIHFTFPSSGFHNDPNGLVY--TG
S. occidentalis   11 : PLTTTFGGYVASSSIDLSVDTSEYNRPLIHFTPEKGHMNDPNGTFYDKTA
S. cerevisiae SUC2  3 : LQAFFTLAGFAAKHSASHTNETSDRPLVHFTFPKGMNDPNGLWIDEKD
                      ..  ....  *  ....  .

S. pombe inv1      101 : GVYHMFQYSPKILTAGE-VHHGHTVSKDLIHWENYPIAIYPDEHENGVL
S. occidentalis    61 : KTWHLTFQYNPNATAHGQPLYHGHATSNDLVHWDHEMAIGPEHNEGIF
S. cerevisiae SUC2  53 : AKWHTYFQYNPNOTVWGTPLFWGHATSDDLTNWEOQPIAIAPKRNDG--
                      *  ..  .  *  ....  *  .  .

S. pombe inv1      151 : SLPFSGSIVVDVHN-----SSGLFSNDTI-PEERIVLIYTDHWTGVAERO
S. occidentalis    111 : ----SGSIVVDHNN-----TSGFF-NSSIDPNQRIVAIYTNHMPDL-QTO
S. cerevisiae SUC2  103 : --AFSGSMVVDYNN-----TSGFF-NDTIDPRQRCVAIWT-YNTPESSEEQ
                      ....  *  .  .  *  .  .  .  .

S. pombe inv1      201 : AIAITTDGGYTFKYSGNPVLIDNSLQFRDPKVMHDFDANRWVIVAMSQ
S. occidentalis    161 : DIAFSLDCCYTFTRYENNPVIDVSSNQFRDPKVFHHERFKSHDHGCSEIA
S. cerevisiae SUC2  153 : YISYSTDGGYTFTEYQRNPVLAANSTQFRDPKVFHYEP6QAWIMTAARSQ
                      *  ....  .  .  .  .  .  .  .


S. pombe inv1      251 : NYGIAFYSSYDLIHHTLSVVFSTSGYLGCLQYECPCGHARVPVECTDEYK-W
S. occidentalis    211 : RVKIQIFG9ANLKNHVLNSNFS-SGYGNQYGH9RLIEVPIENS9DK9K-W
S. cerevisiae SUC2  203 : DYKIEIYSSDDLK9HRT9SAFANE9CL9Y9EC9CLIE9VTE9Q9PSK9YH
                      *  .  .  .  .  .  .  .  .

S. pombe inv1      301 : VLFISINPGAP-LGGSVVQYFVGDNWGTN9FVPD9QTRFVDL9KDFYA-S
S. occidentalis    261 : VMFLAINPGSP-LGGSINQYFVGDFDGFQFVPD9SQTRFVDIGKDFYA-F
S. cerevisiae SUC2  253 : VMFISINPGAP-AGGSFNQYFVG9FN9THF9AF9DNQ99RV9V9FGKDIYALQ
                      *  ....  .  .  .  .  .  .  .

S. pombe inv1      351 : ALIHSS9ANADVIGVGHAS9H9YTN9APTQVF-R9AMTVARKFTLRDVPQ
S. occidentalis    311 : QTF--9EVEHGVLGLAHAS9H9YADQVPTN9H-R9STSLARNYTL9YVH9
S. cerevisiae SUC2  303 : TTFNTDPTYGSALGIAHAS9H9Y9AFVPTN9H-R99HSLVRK9SLNT9YQ
                      ....  .  .  .  .  .  .

```

FIG. 2



F I G . 3





FIG. 4

(a)



(b)





FIG. 5

(a)



(b)






FIG. 6





F I G . 7



# FIG. 8


invertase activity (U/OD)

repression   depression

wild type   1   31

$\Delta inv1+$    0   0

1.0 kb



$\Delta (-2810/-2810)$    2   23



$\Delta (-2810/-2190)$    47   40



$\Delta (-2810/-1710)$    46   38



$\Delta (-2810/-1200)$    51   42



$\Delta (-2810/-200)$    1   1



-2000   -1000   0   1000



FIG. 9






FIG. 10




# FIG. 11



# FIG. 12

(a)



(b)






FIG. 13



## FIG. 14










FIG. 15



F I G . 16





Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) EP 1 223 219 A3

(12)

## EUROPEAN PATENT APPLICATION

(88) Date of publication A3:  
23.10.2002 Bulletin 2002/43

(43) Date of publication A2:  
17.07.2002 Bulletin 2002/29

(21) Application number: 02004252.9

(22) Date of filing: 30.10.1998

(51) Int Cl.7: C12N 15/31, C07K 14/39,  
C12N 15/81, C12N 1/19,  
C12P 21/02  
// (C12N15/31, C12R1:645),  
(C12N1/19, C12R1:645)

(84) Designated Contracting States:  
DE FR GB

(30) Priority: 31.10.1997 JP 31460897

(62) Document number(s) of the earlier application(s) in  
accordance with Art. 76 EPC:  
98950470.9 / 0 972 834

(71) Applicant: ASAHI GLASS COMPANY LTD.  
Tokyo 100-8405 (JP)

(72) Inventors:

- Tohda, Hideki, Asahi Glass Company Ltd.  
Yokohama-shi, Kanagawa 221-0863 (JP)
- Hama, Yuko, Asahi Glass Company Ltd.  
Yokohama-shi, Kanagawa 221-0863 (JP)
- Kumagai, Hiromichi, Asahi Glass Company Ltd.  
Yokohama-shi, Kanagawa 221-0863 (JP)

(74) Representative: Müller-Boré & Partner  
Patentanwälte  
Grafinger Strasse 2  
81671 München (DE)

(54) Inducible promoter and secretion signal for use in *Schizosaccharomyces pombe*, expression vector containing them and their use

(57) The present invention relates to an inducible promoter gene for use in the fission yeast *Schizosaccharomyces pombe*, an inducible expression vector containing the inducible promoter gene, *Schizosaccharomyces pombe* transformed with the inducible expression vector and a process for producing a heterologous protein by using the *Schizosaccharomyces pombe* transformant.

The present invention also relates to a secretion signal gene for use in *Schizosaccharomyces pombe*, an expression vector containing the secretion signal, *Schizosaccharomyces pombe* transformed with the expression vector and a process for producing a heterologous protein by using the *Schizosaccharomyces pombe* transformant. These inducible promoter gene and secretion signal gene are originated from the *S. pombe* gene involved in expression of the invertase precursor.

The inducible promoter gene is originated from the region responsible for the catabolite repression of the

invertase precursor. When a *Schizosaccharomyces pombe* transformant carrying the inducible expression vector is incubated, the timing of the production of the desired heterologous protein can be controlled by regulating the expression of the heterologous protein structural gene depending on the presence or absence of a specific nutrient (especially glucose).

The secretion signal gene encodes the secretion signal for the invertase precursor. When a *Schizosaccharomyces pombe* transformant carrying an expression vector containing the secretion signal gene is incubated, a heterologous protein fused with the secretion signal is synthesized as a precursory protein in the cell, and after intracellular separation of the secretion signal, the heterologous protein is secreted from the cell. Therefore, it is possible to obtain the desired protein from the cell culture, and the desired heterologous protein can be purified easily.

EP 1 223 219 A3



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 02 00 4252

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X	MORENO S ET AL: "PURIFICATION AND CHARACTERIZATION OF THE INVERTASE FROM SCHIZOSACCHAROMYCES POMBE" BIOCHEMICAL JOURNAL, PORTLAND PRESS, LONDON, GB, vol. 267, 1990, pages 697-702, XP002917397 ISSN: 0264-6021 * the whole document *	1-7	C12N15/31 C07K14/39 C12N15/81 C12N1/19 C12P21/02 //(C12N15/31, C12R1:645), (C12N1/19, C12R1:645)
X	DATABASE EMBL 'Online! 3 January 1997 (1997-01-03) YOSHIOKA, S.: "Schizosaccharomyces pombe mRNA, partial cds, clone: SY 1543" retrieved from EBI Database accession no. SPD242 XP002169853 * abstract *	1-7	
P, X	TANAKA N ET AL: "ISOLATION AND CHARACTERIZATION OF AN INVERTASE AND ITS REPRESSOR GENES FROM SCHIZOSACCHAROMYCES POMBE" BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, ACADEMIC PRESS INC. ORLANDO, FL, US, vol. 245, no. 1, April 1998 (1998-04), pages 246-253, XP002917398 ISSN: 0006-291X * page 246, right-hand column, last paragraph * * figure 1 *	1-7	TECHNICAL FIELDS SEARCHED (Int.Cl.7)  C12N C07K
The present search report has been drawn up for all claims			
Place of search <b>THE HAGUE</b>		Date of completion of the search <b>28 August 2002</b>	Examiner <b>Schmitz, T</b>
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons &amp; : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03 92 (P04001)



European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number  
EP 02 00 4252

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
A	NISHIZAWA M ET AL: "CONSTRUCTION OF AN EXPRESSION AND SECRETION VECTOR FOR THE YEAST SACCHAROMYCES-CEREVISIAE" AGRICULTURAL AND BIOLOGICAL CHEMISTRY, vol. 51, no. 2, 1987, pages 515-521, XP002168488 ISSN: 0002-1369 * page 517, right-hand column, last paragraph - page 518, left-hand column, paragraph 2 * * figure 1 * * page 517, left-hand column, paragraph 3 - right-hand column, paragraph 2 *		
A	PEREZ J A ET AL: "CLONING AND SEQUENCE ANALYSIS OF THE INVERTASE GENE INV1 FROM THE YEAST PICHIA ANOMALA" CURRENT GENETICS, NEW YORK, NY, US, vol. 29, 1996, pages 234-240, XP002917395 ISSN: 0172-8083 * the whole document *		
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
Place of search <b>THE HAGUE</b>		Date of completion of the search <b>28 August 2002</b>	Examiner <b>Schmitz, T</b>
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application I : document cited for other reasons &amp; : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03 92 (F04C01)

**THIS PAGE BLANK (USPTO)**